

Request for Renewal of Marine Mammal Protection Act
Incidental Harassment Authorization
NMFS No. 14426

Russian River Estuary Management Project

Applicant:
Sonoma County Water Agency
P.O. Box 11628
Santa Rosa, CA 95406


February 2011

(1) A detailed description of the specific activity or class of activities that can be expected to result in incidental taking of marine mammals.


The Russian River estuary (Estuary) is located about 97 kilometers (km; 60 miles) northwest of San Francisco in Jenner, Sonoma County, California (Figure 1). The Russian River watershed encompasses 3,847 square kilometers (km) (1,485 square miles) in Sonoma, Mendocino, and Lake counties. The Estuary extends from the mouth of the Russian River upstream approximately 10 to 11 km (6 to 7 miles) between Austin Creek and the community of Duncans Mills (Heckel 1994).

The Estuary may close throughout the year as a result of a barrier beach forming across the mouth of the Russian River. The mouth is located at Goat Rock State Beach (California Department of Parks and Recreation). Although closures may occur at anytime of the year, the mouth usually closes during the spring, summer, and fall (Heckel 1994; Merritt Smith Consulting 1997, 1998, 1999, 2000; Sonoma County Water Agency and Merritt Smith Consulting 2001). Closures result in ponding of the Russian River behind the barrier beach and, as water surface levels rise in the Estuary, flooding may occur. Natural breaching events occur when Estuary water surface levels exceed the capability of the barrier beach to impound water, causing localized erosion of the barrier beach and creation of a tidal channel that reconnects the Russian River to the Pacific Ocean.

The barrier beach has also been artificially breached for decades; first by local citizens, then the County of Sonoma Public Works Department, and, since 1995, by the Sonoma County Water Agency (Water Agency). The Water Agency's artificial breaching activities are conducted in accordance with the Russian River Estuary Management Plan recommended in the Russian River Estuary Study 1992-1993 (Heckel 1994). The study was completed to evaluate the impacts of artificial breaching and to select a preferred estuary management program. The purpose of artificially breaching the barrier beach is to alleviate potential flooding of low-lying properties along the estuary. The Water Agency accesses the beach from the paved parking lot at Goat Rock State Beach, located at the end of Goat Rock Road off of Highway 1 (Figure 2). Equipment (e.g a bulldozer, excavator, or similar equipment) is off-loaded in the parking lot and driven onto the beach via an existing access point. A pilot channel in the sandbar is created at a sufficient depth to allow river flows to scour open the mouth of the river and reduce the water surface elevations in the Estuary. As the channel is dug, it first remains disconnected from the Estuary by maintaining a portion of the barrier beach on the Estuary side (as opposed to the ocean side of the beach) to avoid flowing water in the channel. The sand excavated is placed onto the beach adjacent to the pilot channel. After the pilot channel is dug, the last portion of the sandbar adjacent to the Estuary is removed, allowing river water to flow to the ocean. The size of the pilot channel varies depending on the height of the barrier beach to be breached, the tide level, and the water surface elevation in the Estuary. A typical pilot channel would be approximately 100 feet long, 25 feet wide, and 6 feet deep. The amount of sand moved can range from less than 100 cubic yards to approximately 1,000 cubic yards. After the last portion of the barrier beach is removed, water begins flowing out of the channel, scouring and



\\SD-DATA\Pro\Special_projects\Russian_river\ESTUARY-Management-MMPA-IHA.dwg March 1, 2011



Purpose: Russian River Estuary Breaching
Datum: 1929 MSL
Property Owners Adjacent to Project:

SONOMA COUNTY WATER AGENCY
404 Aviation Boulevard
Santa Rosa, CA. 95403

In: Russian River
At: Jenner
County Of: Sonoma, CA.
Application By: SCWA

Figure 2

enlarging the channel to widths of 50 to 100 feet within one or two tidal cycles. Very rapid enlargement, from approximately 25 feet to over 200 feet, has been observed (Heckel 1994).

From 1996 to 2010, the barrier beach was breached during every month of the year, but the majority of breaching events occurred in the fall (October and November), followed by the spring (April, May, and June) and the month of September (Figure 3). The number of artificial breaching events varies each year (Table 1). The lowest number of breaching events occurred in 2004 (1 event) and the highest number was 15 attempted breaches (13 successful artificial breaching events) in 2009. It is difficult to predict how many artificial breaching events are required each year, but there have been an average of 6 artificial breaching events annually over the last 14 years.

BIOLOGICAL OPINION AND THE ESTUARY

The Water Agency and the U.S. Army Corps of Engineers (Corps) consulted with the National Marine Fisheries Service (NMFS) under Section 7 of the Endangered Species Act (ESA) regarding the potential effects of their operations and maintenance activities, including the Water Agency's estuary management program, on federally-listed steelhead (*Oncorhynchus mykiss*), coho salmon (*O. kisutch*), and Chinook salmon (*O. tshawytscha*). As a result of this consultation, the NMFS issued the Russian River Biological Opinion (NMFS 2008) finding that artificially elevated inflows to the Russian River estuary during the low flow season (May through October) and historic artificial breaching practices have significant adverse effects on the Russian River's estuarine rearing habitat for steelhead, coho salmon, and Chinook salmon. The historic method of artificial sandbar breaching, which is done in response to rising water levels behind the barrier beach, adversely affects the estuary's water quality and depths.

The historic artificial breaching practices create a tidal marine environment with shallow freshwater depths and high salinity. Salinity stratification contributes to low dissolved oxygen at the bottom in some areas. The Biological Opinion (NMFS 2008) concludes that the combination of high inflows and breaching practices impact rearing habitat because they interfere with natural processes that cause a freshwater lagoon to form behind the barrier beach. Fresh or brackish water lagoons at the mouths of many streams in central and southern California often provide depths and water quality that are highly favorable to the survival of rearing salmon and steelhead.

The Biological Opinion's Reasonable and Prudent Alternative (RPA) 2 (NMFS 2008) requires the Water Agency to collaborate with NMFS and to modify estuary water level management in order to reduce marine influence (high salinity and tidal inflow) and promote a higher water surface elevation in the estuary (formation of a fresh or brackish lagoon) for purposes of enhancing the quality of rearing habitat for juvenile (age 0+ and 1+) steelhead from May 15th to October 15th (referred to hereafter as the "lagoon management period").¹ A program of

¹ The lagoon management period is May 15th to October 15th, as described in the Russian River Biological Opinion (NMFS 2008).




Figure 3. The number of Russian River estuary breaching events from 1996 to 2010.

Table 1. Breaching of the Russian River Estuary from 1996 to 2010. Number of times breached by year and month, including artificial breaches by Sonoma County Water Agency, natural breaches (denoted by [#]), and breaches conducted by private individuals without a Corps permit, denoted by (#).

Month	Year															
	1996	1997	1998*	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	
January						1								1, [1]	1	
February												2				
March		1, [1]						[1]								
April		[1]				2			[1]			3	[1]	[1]		
May		1, [1]			1	3			1				5			
June		2		1	1		1	[1]						1		
July	1			1									1		{1},[1]	
August	(2)	1							[1]							
September	1, (1)	2	4	1	1					1			1	1		
October	1	1	3	2	2	2	[1]	2	(1)	1	[1]	[1]	1	4	2,[1]	
November	[1]	1	1	1, [1]	4	[1]	3	1	(2)	2	[3]	2	1	4	[1]	
December					2		1					[1]	2	1, [1]	4	
TOTAL	7	12	8	7	11	9	6	5	6	4	5	10	12	17	6	
SCWA	3	9	8	6	11	8	5	3	1	4	0	9	10	15	3	

* Type of breach was not recorded for 1998. All breaching events for 1998 are treated as done by Sonoma County Water Agency.

potential, incremental steps are prescribed to accomplish this, including adaptive management of a lagoon outlet channel on the barrier beach.

Harbor seals (*Phoca vitulina richardii*) regularly haul out at the mouth of the Russian River (Jenner haulout) (Figure 4). California sea lions (*Zalophus californianus*) and northern elephant seals (*Mirounga angustirostris*) are occasionally observed at the haulout. There are also several known river haulouts at logs and rock piles in the Russian River estuary (Figure 4). The Water Agency is applying for incidental harassment authorization (IHA) under the Marine Mammal Protection Act (MMPA) for activities associated with Russian River Estuary Management Activities. These activities include:

- construction and maintenance of a lagoon outlet channel that would facilitate management of a barrier beach (closed sandbar) at the mouth of the Russian River and creation of a summer lagoon to improve rearing habitat for listed steelhead as mandated by the Russian River Biological Opinion (NMFS 2008);
- artificially breaching the barrier beach to minimize the potential for flooding of low-lying properties along the Estuary; and
- monitoring activities associated with the management actions described above.

Lagoon Outlet Channel Management. To comply with the Russian River Biological Opinion, the Water Agency plans to adaptively manage water surface elevations between May 15th and October 15th (lagoon management period) after a barrier beach forms and creates a lagoon.² Modifications to the barrier beach would be small departures from the existing beach and channel topography at the time of closure, and the new channel would be similar to the channel configurations resulting from previous breaching practices and consistent with natural processes. Material excavated for the construction of the outlet channel would be smoothed and graded to a height of approximately 1 to 2 above existing grade. The outlet channel would be constructed to dimensions which do not significantly depart from channels which have been historically observed at the site.

The adaptive lagoon outlet channel management plan seeks to work with natural processes and site conditions to maintain an outlet channel that reduces tidal inflow of saline water into the estuary (PWA 2010). To avoid tidal inflow and maintain a lagoon system that would not flood properties adjacent to the Estuary, the Water Agency would create and maintain a shallow, “perched” outlet channel that would not be excavated as deeply, narrowly, or with as steep a gradient as typical artificial breaching pilot channels, which are designed to allow the current velocities to erode a wider and deeper channel and downcut into the barrier beach.

Active management of estuarine/lagoon water levels would commence when oceanside wave action pushes sand landward to form a natural barrier beach across the river’s mouth. When this happens, the Water Agency would monitor lagoon water surface elevation, as river inflow

² The Russian River Biological Opinion (NMFS 2008) establishes the lagoon outlet channel management period as May 15th to October 15th.



DISCLAIMER

This map document and associated data are distributed for informational purposes only "AS-IS" at the published scale and provided without warranty of any kind expressed or implied. The positional accuracy of the data is approximate and not intended to represent survey map accuracy. The Sonoma County Water Agency assumes no responsibility arising from use of this information.

Pinniped Haulouts at the Russian River Estuary



Figure 4

to the newly closed lagoon builds up behind the barrier beach, causing water surface elevation to rise in the lagoon. The goal is to manage lagoon water surface elevations between 4 and 9 ft National Geodetic Vertical Datum (NGVD)³, which is high enough to enhance fish habitat (NMFS 2008) while also minimizing flood hazard to low-lying structures adjacent to the Estuary (Heckel 1994).

The outlet channel would be excavated and maintained with one or two pieces of heavy machinery (e.g., excavator or bulldozer). The outlet channel would be excavated with a bed elevation 0.5 to 2.0 ft below the lagoon water surface elevation along its entire length to allow outflow from the lagoon to pass over the sandbar. The outlet channel dimensions are estimated to be approximately 30-feet wide, based on a wide and short channel alignment that would minimize scour potential and potential constraints of the acceptable excavation volumes under regulatory permits. The outlet channel would be cut into the top of the naturally formed barrier beach. The outlet channel bed slope would be minimized to reduce the potential for bed scour and unintentional breaching of the sandbar. The outlet channel width and length estimates are consistent with historic river mouth widths and lengths observed within the lagoon management period (Behrens 2008).

The channel's length is estimated to vary from 100 to 400 ft, consistent with historic channel lengths observed within the management period (Behrens 2008). Length would be a function of the channel's planform alignment⁴. Planform alignment of the channel would vary within the region in which the channel has been observed to naturally occur (Figure 2). The southern extent of this region would be the jetty and would extend approximately 1,500 feet to the northwest. Various channel locations may be pursued in an effort to adapt other project variables, such as bed slope, bed elevation and channel width, and to take advantage of site features such as areas of reduced wave energy. For example, alignment at the start of the management period may follow the northward alignment typically observed at this time of year to take advantage of the low berm crest elevation along this alignment. However, the channel may migrate from this initial alignment. If the channel then closes, alternative channel alignments within the region shown in Figure 2 may be implemented to test the relationship of mouth location on channel stability.

The strategy for outlet channel configuration and modifications would be an incremental approach that seeks to minimize the risk of uncontrolled breaching which returns the estuary to tidal conditions. The precise number of outlet channel implementation events would depend on uncontrollable variables such as seasonal ocean wave conditions (e.g. wave heights and lengths), river inflows, and the success of previous excavations (e.g. the success of selected channel widths and meander patterns) in forming an outlet channel that effectively maintains lagoon water surface elevations. It is predicted that up to three successive outlet channel excavations, at increasingly higher beach elevations, may be necessary, with the result being a "perched" lagoon. The goal is to develop an outlet channel that supports a stable "perched"

³ Water surface elevations are measured by the Agency's gage located at the State Parks Visitor Center in Jenner.

⁴ Planform alignment is the centerline alignment of the channel in planimetric view.

lagoon with water surface elevations at approximately 7 ft NGVD for several months. Stable conditions imply that river inflow into the lagoon would be approximately the same as outflow through the outlet channel and that net sand deposition or erosion does not impair the outlet channel's function.

In the event that a "perched" outlet channel fails (*i.e.*, erodes the barrier beach and forms a tidal inlet), the Water Agency would resume adaptive management of the outlet channel's width, slope, and alignment in consultation with the NMFS and California Department of Fish and Game (CDFG) after ocean wave action naturally reforms a barrier beach and closes the river's mouth during the lagoon management period.

Additional details regarding the development of the adaptive lagoon outlet channel management plan may be found in PWA (2010) attached to this application.

Implementation and Maintenance. The Water Agency would contact State Parks lifeguards, as well as State Park District headquarters and the Monte Rio Fire Protection District, within 24 hours prior to excavating and maintaining the lagoon outlet channel to minimize potential hazards to beach visitors. Signs and barriers would be posted 750 feet of each side of the outlet channel for 24 hours prior to and after excavation events to warn beach visitors of the hazards of the area and the presence of pinnipeds on the beach. Notifications for the general public would also be posted at the Jenner visitor's center boat launch.

The barrier beach would be accessed from the paved parking lot at Goat Rock State Beach, located at the end of Goat Rock Road off of Highway 1 (Figure 2). Equipment would be off-loaded in the parking lot and driven north onto the beach via an existing access point. Water Agency crews would approach the haulout ahead of the heavy equipment to minimize the potential for flushes to result in a stampede, a particular concern during pupping season. Water Agency staff would avoid walking or driving equipment through the seal haulout. Crews on foot would take caution to approach the haulout slowly and to make an effort to be seen from a distance, if possible, rather than appearing suddenly at the top of the sandbar. Personnel on the beach would include up to two equipment operators, three safety team members on the beach (one on each side of the channel observing the equipment operators, and one at the barrier to warn beach visitors away from the activities), and one safety team member at the overlook on Highway 1 above the beach. Occasionally, there would be two or more additional people on the beach (Water Agency staff or regulatory agency staff) on the beach to observe the activities. Water Agency staff would be followed by the equipment, which would then be followed by an Water Agency vehicle (typically a small pickup truck, the vehicle would be parked at the previously posted signs and barriers on the south side of the excavation location). Equipment would be driven slowly on the beach and care would be taken to minimize the number of shut downs and start ups when the equipment is on the beach.

Creating and maintaining the outlet channel would probably employ one or two pieces of heavy equipment (e.g. excavator or bulldozer) to move sand on the beach. At the start of the management period (late spring or early summer), when configuring the outlet channel for the

first time that year, machinery may operate on up to 2 consecutive working days. As technical staff and maintenance crews gain more experience with implementing the outlet channel and observing its response, it may be possible to reduce the frequency of maintenance during the remainder of the management season. In consideration of the beach environment, effort would be made to minimize the amount and frequency of mechanical intervention, thereby reducing disturbances to seals and other wildlife, as well as State Park's visitors on the beach.

The quantity of sand moved would depend on antecedent beach topography. Excavation volumes would not exceed 2,000 cubic yards. Any sand excavated from the channel would be immediately placed on the adjacent beach within the wave wash zone to promote natural removal to minimize changes to beach topography outside the outlet channel.

The Water Agency anticipates that lagoon outlet channel management activities would occur in accordance with the Russian River Biological Opinion and that they would primarily occur between May 15th and October 15. However, if estuary water surface elevations rise above 7.0 feet (at the Jenner gage) during the lagoon management period, the Water Agency may consult with NMFS and CDFG regarding artificially breaching the sandbar to alleviate potential flooding, as discussed in the Biological Opinion and described below.

ARTIFICIAL BREACHING

Artificial breaching activities occur on the closed sandbar. The Water Agency mechanically breaches the sandbar to alleviate potential flooding of low-lying shoreline properties near the town of Jenner. For more than a decade, breaching has been performed in accordance with the *Russian River Estuary Study 1992-1993* (Heckel 1994) when the Estuary water surface level is between 4.5 and 7.0 feet as read at the Jenner gage (located at the Jenner Visitor's Center). The Water Agency would contact State Parks lifeguards, as well as State Park District headquarters and the Monte Rio Fire Protection District, within 24 hours prior to breaching activities to minimize potential hazards to beach visitors. Signs and barriers would be posted 750 feet of each side of the pilot channel for 24 hours prior to and after breaching events to warn beach visitors of the hazards of the breaching area and the presence of pinnipeds on the beach. Notifications for the general public would also be posted at the Jenner visitor's center boat launch.

The barrier beach would be accessed from the paved parking lot at Goat Rock State Beach, located at the end of Goat Rock Road off of Highway 1 (Figure 2). Equipment would be off-loaded in the parking lot and driven north onto the beach via an existing access point. Water Agency crews would approach the haulout ahead of the heavy equipment to minimize the potential for flushes to result in a stampede, a particular concern during pupping season. Water Agency staff would avoid walking or driving equipment through the seal haulout. Crews on foot would take caution to approach the haulout slowly and to make an effort to be seen from a distance, if possible, rather than appearing suddenly at the top of the sandbar. Personnel on the beach would include an equipment operator, three safety team members on the beach (one on each side of the channel observing the equipment operators, and one at the barrier to warn beach visitors away from the breaching activities), and one safety team member

at the overlook on Highway 1 above the beach. Occasionally, there would be two or more additional people on the beach (Water Agency staff or regulatory agency staff) on the beach to observe breaching activities. Water Agency staff would be followed by the equipment, which would then be followed by an Water Agency vehicle (typically a small pickup truck, the vehicle would be parked at the previously posted signs and barriers on the south side of the excavation location). Equipment would be driven slowly on the beach and care would be taken to minimize the number of shut downs and start ups when the equipment is on the beach. Creating and maintaining the outlet channel would probably employ one excavator or bulldozer) to move sand on the beach.

Breaching activities would typically be conducted on outgoing tides to maximize the elevation head difference between the estuary water surface and the ocean. A cut in the barrier beach would be created at a sufficient depth to allow river flows to begin transporting sand to the ocean. The sand would be placed onto the beach adjacent to the pilot channel. After the pilot channel is dug, the last upstream portion of the sandbar would be removed, allowing river water to flow to the ocean. The size of the pilot channel varies depending on the height of the sandbar to be breached, the tide level, and the water surface elevation in the Estuary. A typical channel would be approximately 100 feet long, 25 feet wide, and 6 feet deep. The amount of sand moved would range from less than 100 cubic yards to approximately 2,000 cubic yards.

The Water Agency anticipates that artificial breaching activities would occur in accordance with the Russian River Biological Opinion and that they would primarily occur from October 16th to May 14. However, if estuary water surface elevations rise above 7.0 feet (at the Jenner gage) during the lagoon management period (May 15th through October 15th), the Water Agency would artificially breach the sandbar to alleviate potential flooding, as discussed in the Biological Opinion. The Biological Opinion incidental take statement estimates that the Water Agency may need to artificially breach the sandbar “twice per year between May 15 and October 15 during the first three years covered by this opinion, and once per year between May 15 and October 15 during years 4-15 covered by this opinion” (NMFS 2008).

MONITORING

Implementation of the lagoon outlet channel adaptive management plan would require monitoring to measure changes in the bar and channel elevation, lengths, and widths, as well as flow velocities and observations of the bed structure (to identify bed forms and depth-dependent grain size distribution indicative of armoring) in the channel. In addition to the activities described for the lagoon outlet channel adaptive management plan, the Water Agency is required by the Russian River Biological Opinion and other state and federal permits to collect biological, water quality, and physical habitat data in conjunction with estuary management. Fisheries seining and trapping, water quality monitoring, invertebrate/sediment sampling, and physical habitat measurements require the use of boats and nets in the Estuary. Boating and other monitoring activities occur in the vicinity of river haul outs (see Figure 4, Mortenson 2009). Table 2 provides a summary of the monitoring tasks and the frequency of their implementation.

Table 2. Monitoring tasks associated with Russian River estuary management with potential to disturb pinnipeds.

Task	Description	Field Activities	Frequency
Lagoon Outlet Channel Management on the Barrier Beach			
Discharge Measurements	Collected within the outlet channel to verify the channel's conveyance.	2 field staff to complete cross sectional flow velocity surveys using flow meter attached to a wading rod with electronic data logger (beeps); bank pins to be installed on either bank and fiberglass measuring tape stretched from bank to bank.	Every 2 weeks
Outlet Channel Bed Structure	Observe the bed for bed forms and depth-dependent grain size distribution indicative of armoring. Sediment sampler used.	2 field staff to collect sediment sample from the surface of the channel bed.	Monthly
Outlet channel topography	Collect outlet channel elevation and width	2 field staff would capture outlet channel features using a prism mounted on a survey rod.	Monthly
Biological and Physical Habitat Monitoring in the Estuary			
Fisheries seining	Deploy seine to collect fish at up to 8 locations in the estuary	One or two boats with approximately 6 field staff	Monthly
Invertebrate/salmonid prey study	Collection of benthic invertebrates and zooplankton	One boat with 2 field staff	Weekly
Water quality	Collection of temperature, dissolved oxygen, conductivity, pH, depth, nutrient and bacteriological samples	A boat with 2 or 3 field staff, 6 datasonde arrays submerged in estuary at various locations from mouth to Duncans Mills.	Every 3 weeks
SCWA topographic survey of sandbar	Survey of sandbar height and widths	2 field staff on beach equipped with a survey rod.	Monthly
BML flow circulation (under contract w/SCWA)	Survey of cross sectional velocity data in estuary and collection of temperature and salinity profile data at various locations from mouth to Duncans Mills.	A boat with 2 or 3 field staff, collecting cross sectional data from mouth to Duncans Mills.	Weekly

(2) The date(s) and duration of such activity and the specific geographical region where it would occur.

The anticipated marine mammal disturbance from project activity would occur at the mouth of the Russian River (38.450833, -123.129873) in Jenner, California. The Russian River estuary is located about 97 km (60 miles) northwest of San Francisco. The harbor seals primarily haul out on the estuary-side of the beach (Figures 2 and 4).

The Estuary closes throughout the year as a result of a sandbar forming at the mouth of the Russian River. To facilitate summer lagoon management, the Water Agency would construct the lagoon outlet channel after the first natural barrier beach closure, but the lagoon would generally be managed from May 15th to October 15th (Table 3). It is anticipated that the outlet channel implementation would be a 2-day event with initial construction of the lagoon outlet channel taking one day of work, and subsequent adjustments to the outlet channel on the second day. Subsequent maintenance would occur approximately weekly through October 15th. Artificial breaching activities would generally occur between October 16th and May 14th (Table 3). Biological and water quality monitoring generally occurs from mid-April through December (Table 2).

(3) The species and numbers of marine mammals likely to be found within the activity area.

The species of marine mammals that are likely to occur in the project area include the following pinnipeds: harbor seals and the California sea lions. Sightings of sea lions have been reported during the months from December to June, likely foraging, but their numbers are normally low (Hanson 1993). In the last several years, a single male northern elephant seal (*Mirounga angustirostris*) has been present at the Jenner haul out during the late winter and spring.

The number of harbor seals at the Russian River varies throughout the year (Table 4). These numbers have been recorded extensively since 1972 at the mouth of the Russian River, where several local residents, working independently or under the guidance of the Stewards of the Coast and Redwoods, have recorded the harbor seal population at the mouth and within the Russian River. It is believed that harbor seals established the haul out site at the Russian River in 1972 (*i.e.*, the first known records) and their numbers at the site have steadily grown (Hanan and Beeson 1994, Mortenson and Twohy 1994). No estimates of pup numbers during pupping season are known as monitors are unable to distinguish pups from the other age classes from their observation site (J. Mortenson and N. Jellison, pers comm. 2009), but it is assumed that some pupping occurs there. During the months from September to November, the number of seals hauling out on the beach declines significantly. In most cases, harbor seals are not present at the mouth of the Russian River during the fall or if they are present, the numbers are fewer than 50 during this period. The harbor seals normally return in greater numbers during the late winter (February and March) or early spring (April), and remain at the beach in great numbers until the end of July. Although the number of harbor seals at this out has fluctuated from year to year, average counts show a steady rise in population trend. During recent state censuses, the number of harbor seals observed during the single-day summer counts has continued to steadily increase, with nearly 350 seals observed in 1993 (Mortenson and Twohy 1994) and 315 in 2004 (Lowry et al. 2005), although over 500 animals have been recorded (Mortenson and Twohy 1994). Table 4 reflects the monthly mean number of harbor seals recorded by E. Twohy during daily counts of seals at the Jenner haulout from 1993 to 2005 (without differentiating between bar-open and bar-closed conditions). Figure 5 shows the mean number of harbor seals observed at the Jenner haulout during baseline monitoring for the Estuary Management Project from April to December 2010. Table 5 shows the average number of harbor seals observed at the Jenner haulout (Goat Rock State Beach) during bar-

Table 3. Estimated frequency and duration of Russian River estuary management activities with potential to disturb pinnipeds.

Task and Dates	Duration and Frequency	Potential No. of Take Events ^a
Lagoon Outlet Channel Management on the Sandbar (May 15 to October 15)		
Excavation of outlet channel	Daily up to 2 consecutive days per event; up to 3 events estimated.	3
Maintenance of outlet channel	1 day per week	May-1; June-4; July-4; Aug-4; Sept-4; Oct-1 (18 total)
Outlet channel discharge & bed structure measurements	Discharge: ½ day every 2 weeks Structure: ½ day per month (taken on same day as discharge measurements)	10 ^b
Outlet channel topography	1 day per month	
Artificial Breaching on the Sandbar (October 16 to May 14)		
October	Averages ½ day per breaching event	2 ^c
November		2
December		2
January		1
February		1
March		1
April		1
May		1
		11 events maximum
Biological and Physical Habitat Monitoring in the Estuary		
Fisheries seining	6 days, monthly from May to October	6 ^d
Invertebrate/salmonid prey study	Weekly from July to October	14 ^d
Water quality	3 days, every 3 weeks from May to December	11 ^d
SCWA topographic survey of sandbar	1 per month for duration of IHA, averages ½ day	12
BML flow circulation (under contract w/SCWA)	Weekly from May to October	22

^a For implementation of the lagoon outlet channel, an event is defined as a single, 2-day episode. It is assumed that the same individual seals would be hauled out during a single event. For the remaining activities, an event is defined as a single day on which an activity occurs. Some events may include multiple activities.

^b The lagoon outlet channel discharge, bed structure, and channel topography monitoring would occur on the same day each month. The outlet channel discharge is collected every 2 weeks and would require an additional ½ day of work.

^c The number of events is the monthly average number of artificial breaching events from 1996 to 2010 (Table 1). Based on the number of events, the maximum number of artificial breaching events would be 15. The average number of breaching events from 1996 to 2010 is 6 events/year. The number of events was reduced from the data in Table 1 to 1 for the month of April and 1 for the month of May to anticipate the request from NMFS that only a single closure in each of these months be breached if it seemed unlikely that additional closures would occur prior to May 15th. The potential for two closures in the first two weeks of May seems low. The number of artificial breaching events in February was reduced to one because closures in that month occurred only in 2007, which was an unusual circumstance. December was increased from an average of 1 to 2 events because of the recent experience of multiple closures during heavy storm events.

^d Assumption is that pinnipeds may be encountered once per event and flush from river haulout in the Estuary.

Table 4. Average daily number of seals observed at Goat Rock State Beach (near Russian River mouth), for each month from 1993 to 2005. Adapted from Mortenson and Twohy 1994 and Elinor Twohy unpublished data.

Year	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
1993	140	219	269	210	203	238	197	34	8	38	78	163
1994	138	221	243	213	208	212	246	98	26	31	101	162
1995	133	270	254	261	222	182	216	74	37	24	38	148
1996	144	175	261	247	157	104	142	65	17	29	76	139
1997	154	177	209	188	154	119	186	58	20	29	30	112
1998	119	151	192	93	170	213	232	53	33	21	93	147
1999	161	170	215	210	202	128	216	98	57	20	74	123
2000	151	185	240	180	158	245	256	63	46	50	86	127
2001	155	189	161	168	135	212	275	75	64	20	127	185
2002	117	12	20	154	134	213	215	89	43	26	73	126
2003	--	1	26	161	164	222	282	100	43	51	109	116
2004	2	5	39	180	202	318	307	35	40	47	68	61
2005	0	7	42	222	220	233	320	145	--	--	--	--

Months represented by "--" indicate periods where data were missing or incomplete.




Figure 5. Mean number of harbor seals observed at the Jenner haulout (Goat Rock State Beach) during Russian River Estuary Management Project baseline pinniped monitoring from April to December 2010. Pups are counted separately through June, after which all seals are counted as adults as it becomes more difficult to accurately age individuals.

Table 5. Average number of harbor seals observed at the Jenner haulout (Goat Rock State Beach) during bar-closed conditions by month during monitoring of artificial breaching activities from 1996 to 2000. From Merritt Smith Consulting (1997, 1998, 1999 and 2001) and Sonoma County Water Agency and Merritt Smith Consulting (2001).

April	May	June	July	August	September	October	November
173	103	100	75	17	5	22	11

closed conditions by month during monitoring of artificial breaching activities from 1996 to 2000. Figure 6 shows the mean number of harbor seals observed at the Jenner haulout (Goat Rock State Beach during barrier beach formed (bar-closed) conditions by month during pinniped monitoring for Russian River Estuary Project water level management activities from April to December 2010.




Figure 6. Mean number of harbor seals observed at the Jenner haulout (Goat Rock State Beach) during pinniped monitoring for Russian River Estuary Project water level management activities from April to December 2010.

(4) A description of the status, distribution, and seasonal distribution (when applicable) of the affected species or stocks of marine mammals likely to be affected by such activities.

Pacific harbor seals-California stock. The California stock of Pacific harbor seals (*Phoca vitulina richardii*) ranges from Cedros Island (Baja California) along the Pacific coasts of the United States, Canada and Alaska, through the Aleutian Islands to the Pribilof Islands. In California, approximately 400-500 harbor seal haul out sites are widely distributed along the mainland and on offshore islands, including intertidal sandbars, rocky shores and beaches (Hanan 1996). California harbor seals are not listed under the ESA or considered strategic under the Marine Mammal Protection Act (MMPA).

The most recent minimum population estimate of the harbor seal population is approximately 31,600 (Carretta *et al.* 2007). Counts of harbor seals in California showed a rapid increase from approximately 1972 (when the MMPA was passed) to 1990. Net production rates appeared to decline from 1982 to 1994. Although earlier analyses were equivocal (Hanan 1996) and there has been no formal determination that the California stock has reached its Optimal Sustainable Population level (defined in the MMPA), the decrease in population growth rate has occurred at the same time as a decrease in human-caused mortality and may be an indication that the population is reaching its environmental carrying capacity.

In general, harbor seals do not undertake long migrations, but do travel 300-500 km on occasion to find food or suitable breeding areas (Herder 1986). Harbor seals are rarely found in pelagic waters and typically stay within the tidal and intertidal zones. On land, harbor seals haul out on rocky outcrops, mudflats, sandbars and sandy beaches with unrestricted access to water and with minimal human presence. Haulout sites are important as resting sites for harbor seals. Harbor seals feed opportunistically in shallow waters on fish, crustaceans, and cephalopods. Foraging occurs in shallow littoral waters, and common prey items include flounder, sole, hake, codfish, sculpin, anchovy and herring (California Department of Fish and Game 2005). Harbor seals are typically solitary while foraging, although small groups have been observed. They normally choose isolated sites for pupping. Pupping normally occurs at the Russian River from March until late June, and sometimes into early July (Mortenson 2009).

Harbor seals have many haulout sites in Northern California with approximately 6 primary mainland haul out sites and possibly a total of 17 haul out sites, if smaller areas are considered, in Sonoma County (Figure 7). The Russian River haul out is the largest in Sonoma County, comprising of approximately 18% of the harbor seal population found there (M. DeAngelis, pers. comm.). There are also several known haulouts in the Russian River estuary at logs and rock outcroppings in the river.

Monitoring efforts are particularly strong in the Point Reyes area, located in Marin County, south of Sonoma County, at the Russian River (Figure 8), and the Gualala River area (south near Sea Ranch). Further north, seals are known to have numerous haul out sites, but monitoring efforts are sparse in the stretch of coastline between the Gualala River area and Humboldt Bay (Figure 9).




Figure 7. Pinniped Haulouts from Bodega Bay to Cuffey's Point, California Coast. Data includes haul-outs and rookeries surveyed using aerial or ground count methodologies and observations and monitoring results. Com_Name is the species common name. *Source:* California Pinniped Rookeries & Haul-out Sites. GIS public map. NOAA National Marine Fisheries Service Southwest Regional Office. <http://www.swr.noaa.gov/psd/rookeryhaulouts/index.htm>




Figure 8. Pinniped Haulouts from Point Arena to Ten Mile River, California Coast. Data includes haul-outs and rookeries surveyed using aerial or ground count methodologies and observations and monitoring results. *Source:* California Pinniped Rookeries & Haul-out Sites. GIS public map. NOAA National Marine Fisheries Service Southwest Regional Office. <http://www.swr.noaa.gov/psd/rookeryhaulouts/index.htm>



Figure 9. Pinniped Haulouts from Caspar Point to Humboldt Bay, California Coast. Data includes haul-outs and rookeries surveyed using aerial or ground count methodologies and observations and monitoring results. *Source:* California Pinniped Rookeries & Haul-out Sites. GIS public map. NOAA National Marine Fisheries Service Southwest Regional Office. <http://www.swr.noaa.gov/psd/rookeryhaulouts/index.htm>

Mortenson (1996) discussed that the number of seals present at the Jenner haulout declined during bar closed (barrier beach closed) conditions. The Water Agency's pinniped monitoring from 1996 to 2000 focused on the barrier beach artificial breaching activities and its effects on the Jenner haulout. Seal counts and disturbances were recorded from 1 to 2 days prior to breaching, the day of breaching, and the day after breaching (Merritt Smith Consulting 1997, 1998, 1999, 2000; Sonoma County Water Agency and Merritt Smith Consulting 2001). In each year, the trend observed was that harbor seal numbers declined during a beach closure (occasionally, the numbers rose again and then declined again during a closure) and increased the day following an artificial breaching event. Observations of disturbances to the Jenner haulout show that the numbers of seals at the haulout (during barrier beach closures) were higher in the morning than later in the day. While seals often alerted to distance sources of disturbance, such as the sound of trucks braking on Highway 1 nearby, seals primarily fled the haulout as a result of disturbances on the beach. The number of seals declined during the day due to disturbances by people on the beach or kayakers/boaters approaching the haulout. Disturbances on the beach typically increased as the morning progressed (greater number of visitors on the beach in the late mornings and early afternoons). There were numerous occasions when the Water Agency's heavy equipment was in operation, but the seals did not leave the haulout or flushed into the water in low numbers.

According to Heckel (1994), "the loss of easy access to the haulout and ready escape to the sea when the river mouth is closed may account for the lower number of harbor seals seen at that time." The mouth of the Russian River is typically open during the winter months, but intermittently closes during the late spring through fall (Figure 2).

Joe Mortenson began his ongoing monthly seal counts at the Jenner haulout and Bodega Rock in January 1987, with nearby haulouts added to the counts thereafter. Elinor Twohy began daily counts of seals and people at the Jenner haulout, including photographing the haulout, on November 1, 1989. Her daily counts were taken at different times on successive days to determine if there were diurnal patterns in use of the haulout (Mortenson and Twohy 1994). She also photographed and noted whether the mouth at the Jenner haulout was opened or closed each day. The information that has emerged from these data sets is that the Jenner haulout is atypical in terms of the time of year that the peak numbers of harbor seals are present. The numbers of seals at the Jenner haulout peaks in the late winter (February and March); at other harbor seal haulouts, peaks are typically observed during the pupping and molting season (spring and summer; Mortenson and Twohy 1993). The Jenner haulout is also atypical in terms of the time of day seal count peaks are observed. At other harbor seal haulouts, daily peaks are typically observed at midafternoon low tides regardless of the season. Although daily harbor seal numbers at the Jenner haulout do peak at midday during the winter (November 16th to March 30th) and in the pupping and molting seasons (April/May and June/July/August, respectively), a midday peak is not observed during the fall (Mortenson and Twohy 1994).⁵

⁵ The winter, pupping, and molting seasons were defined in Mortenson and Twohy 1994).

Mortenson (1996) observed pups were first seen at the Jenner haulout in late March, with maximum counts in May. In this study, pups were not counted separately from other age-classes at the haulout after August due to the difficulty in discriminating pups from small yearlings (Mortenson 1996). Hanson (1993) observed during her study from August 1989 to July 1991 that pupping began at the Jenner haulout in mid-April, with a maximum number of pups observed during the first two weeks of May. This corresponds with the peaks observed at Point Reyes, where the first viable pups are born around the first to second week of March and the peak is the last week of April to early May (Mortenson and Allen, pers. comm.). Pupping season is defined as March 15 to June 30 at the Jenner haulout by the NMFS Incidental Harassment Authorization No. 14426 dated March 31, 2010.

California sea lions-U.S. stock. California sea lions (*Zalophus californianus*) range from southern Mexico to British Columbia, Canada. The entire U.S. population has been estimated at 238,000, and growing at a rate of approximately 6.52% annually between 1975 and 2005 (Carretta *et al.* 2007). The population has experienced an annual growth rate of approximately 6% since at least 1975. The species is not listed under the Endangered Species Act and is not “depleted” or listed as “strategic” stock under the MMPA. Sea lions can be found at sea from the surf zone out to near shore and pelagic waters. On land, the sea lions are found resting and breeding in groups of various sizes, and haul out on rocky surfaces and outcroppings and beaches, as well as manmade structures such as jetties and beaches. Sea lions prefer haul out sites and rookeries near abundant food supplies, with easy access to water; although sea lions occasionally travel up rivers and bays in search of food.

Sea lions exhibit seasonal migration patterns organized around their breeding patterns. California sea lions breed at large rookeries on the Channel Islands in southern California, and on both sides of the Baja California peninsula, typically from May to August. Females tend to remain close to the rookeries throughout the year, while males migrate north after the breeding season in the late summer, and then migrate back south to the breeding grounds in the spring (California Department of Fish and Game 1990). No established rookeries are known north of Point Reyes, California, but large numbers of subadult and non-breeding or post-breeding male California sea lions are found throughout the Pacific Northwest. There is a mean seasonal pattern of peak numbers occurring in the northwest during fall, but local areas show high annual and seasonal variability.

Sea lions feed on fish and cephalopods, including Pacific whiting, rockfish, anchovy, hake, flatfish, small sharks, squid, and octopus (California Department of Fish and Game 1990). Although solitary feeders, sea lions often hunt in groups, which can vary in size according to the abundance of prey (California Department of Fish and Game 1990).

Solitary California sea lions were occasionally observed between the river mouth and the Jenner visitor’s center during bar-open conditions in the Russian River estuary (Merritt Smith Consulting 1999 and 2000). A single sea lion was hauled out during post-breaching monitoring on September 6, 2000 (Sonoma County Water Agency and Merritt Smith Consulting 2001).

Juvenile sea lions were observed during the summer of 2009 at Patty's Rock by Water Agency staff and members of the public. Regular observation of juvenile sea lions were reported along the Sonoma Coast and in the Russian River estuary and were generally considered to be a result of poor foraging conditions in the ocean in 2009. During baseline and artificial breaching monitoring in 2009 and 2010, individual California sea lions were observed at the mouth of the Russian River in November and December 2009. Only a single individual was observed hauling out at the Jenner haulout in November 2009. Other individuals were observed in the surf at the mouth of the river or swimming inside the Estuary. Juvenile California sea lions were observed at Patty's Rock in the Russian River estuary during the summer of 2009 by Water Agency staff and members of the public and sea lions were observed during monitoring of the pinniped peripheral haulouts in October 2009.

Northern elephant seals – California stock. Northern elephant seals (*Mirounga angustirostris*) breed and give birth in California (U.S.) and Baja California (Mexico), primarily on offshore islands (Stewart *et al.* 1994), from December to March (Stewart and Huber 1993). Males feed near the eastern Aleutian Islands and in the Gulf of Alaska, and females feed further south, south of 45°N (Stewart and Huber 1993, Le Boeuf *et al.* 1993). Adults return to land between March and August to molt, with males returning later than females. Adults return to their feeding areas again between their spring/summer molting and their winter breeding seasons. Pups are born in early winter from December to January. Breeding occurs from December to March, and gestation lasts around 11 months. Northern elephant seals are "polygamous"; males establish dominance over large groups of females during the breeding season.

Populations of northern elephant seals in the U.S. and Mexico were all originally derived from a few tens or a few hundreds of individuals surviving in Mexico after being nearly hunted to extinction (Stewart *et al.* 1994). Given the very recent derivation of most rookeries, no genetic differentiation would be expected. Although movement and genetic exchange continues between rookeries, most elephant seals return to their natal rookeries when they start breeding (Huber *et al.* 1991). The California breeding population is now demographically isolated from the Baja California population and is considered to be a separate stock. Based on the estimated 35,549 pups born in California in 2005, the California stock was approximately 124,000 in 2005 (Carretta *et al.* 2009). Based on trends in pup counts, northern elephant seal colonies were continuing to grow in California through 2005 (Carretta *et al.* 2009), but appear to be stable or slowly decreasing in Mexico (Stewart *et al.* 1994).

Northern elephant seals range along the entire California coast (California Department of Fish and Game 2009). Adult male elephant seals breed with harems of females in from mid-December through March in dense rookeries on the San Miguel Island, Santa Barbara Island, San Nicolas Islands, San Simeon Island, Southeast Farallon Island, Año Nuevo Island, on the mainland at Año Nuevo (San Mateo Co.), and the Point Reyes Peninsula (California Department of Fish and Game 2001). From April to November, they feed at sea or haul out to molt at rookeries. They are not listed as "endangered" or "threatened" under the Endangered Species

Act nor as "depleted" or "strategic" under the MMPA. Elephant seals feed at night in deep water, primarily on rays, sharks, pelagic squid, ratfish, and Pacific hake (California Department of Fish and Game 2009). Entanglement in marine debris, fishery interactions, and boat collisions are their main threats.

Censuses of pinnipeds at the mouth of the Russian River have been taken at least semimonthly since 1987. Elephant seals were noted from 1987 to 1991. From 1992-1995, one or two elephant seals were counted during the censuses conducted in May, with occasional records during the fall and winter (Mortenson and Follis 1997). A single male northern elephant seal was present at the mouth of the Russian River harbor seal haul out site, during the late winter and spring for several years. The elephant seal was believed to be a juvenile or sub-adult male when it first began using the area as a haul out site. It was observed harassing harbor seals hauled out at the mouth of the Russian River. Censuses of pinnipeds at the mouth of the Russian River have recorded observations of elephant seals at the mouth of the Russian River in May from 1992 to 1995, with occasional records in the fall and winter (Mortenson and Follis 1997). A northern elephant seal subadult, tagged R-1 by Dr. Sarah Allen in August 2003, was present at the Jenner haulout from 2002 to 2007. He was generally present during molt and again from late December through March into the early pupping season. A single juvenile elephant seal was observed at the Jenner haulout in June 2009.

(5) The type of incidental taking authorization that is being requested (i.e., takes by harassment only; takes by harassment, injury and/or death) and the method of incidental taking.

This is a request for an incidental harassment authorization (IHA, Level B harassment) of harbor seals, California sea lions, and northern elephant seals at the Russian River, in Sonoma County, California. The type of take expected is incidental harassment of pinnipeds from the activities associated with estuary management, which includes people, vehicles, and heavy equipment on the beach near the haulout and activities in the Russian River estuary near river haulout locations. Activities may include: excavation and maintenance of the lagoon outlet channel, construction of a pilot channel during artificial breaching events, posting and removal of warning signs on the beach, monitoring the lagoon outlet channel, monthly topographic surveys of the sandbar at the mouth of the Estuary, boat operation associated with flow circulation and water quality monitoring, and beach-seining and boat operation associated with biological monitoring near haulout locations.

(6) By age, sex, and reproductive condition (if possible), the number of marine mammals (by species) that may be taken by each type of taking identified in paragraph (a)(5) of this section, and the number of times such takings by each type of taking are likely to occur.

The estimates of the number of Pacific harbor seals, California sea lions, and northern elephant seals that may be harassed by the proposed activities is based upon the number of potential

take events associated with Russian River estuary management activities (Table 3) and the average number of individuals of each species that are present at the Jenner haulout during bar-closed conditions (information in response to question 4). The numbers of take events associated with lagoon outlet channel management are split into two categories: 1) initial channel excavation, which would likely occur between May and September, and 2) maintenance and monitoring of the outlet channel, which would continue until October 15th. The Estuary has remained closed for an extended period of time (greater than 14 days) only on a single event (September 2009) since regular counts of pinnipeds at the Jenner haulout began. It is difficult to estimate the numbers of seals that may be hauled out on the barrier beach when the lagoon is formed; however, harbor seals are regularly observed crossing overland from the Pacific Ocean to haul out on the Estuary side of the beach, even in bar-open conditions, so it is anticipated that seals would continue to use the haulout in bar-closed, lagoon conditions. Based on pinniped monitoring from 1996 to 2000 associated with artificial breaching events, the average number of harbor seals hauled out during barrier beach-closed conditions (Table 5) can be used to estimate the number of individuals that may be harassed by both lagoon outlet channel and artificial breaching activities. Both activities would likely be implemented soon after a beach closure (within 14 days), so the data presented in Table 5 would be reasonable for the take estimates from April to November. Because the lagoon outlet channel implementation dates cannot be determined yet (they are dependent on when the barrier beach naturally closes after May 15th), the highest average number of harbor seals presented in Table 5 (May) was used to conservatively estimate the number of seals that may be taken during barrier beach-closed conditions and excavation of the lagoon outlet channel. For maintenance and monitoring activities associated with the lagoon outlet channel, the average number of harbor seals for each month (Table 5) was used. Harbor seal numbers presented in Table 4 were used to estimate take associated with artificial breaching from December to March as this was the best information available for those months and overlapped with the peak in harbor seal numbers at the Jenner haulout. For biological and physical habitat monitoring activities in the Estuary, it was assumed that pinnipeds may be encountered once per event and flush from a river haulout. The estimated potential total number of individual animals that may be taken equates to the maximum number of seals of each species anticipated to be encountered per event multiplied by the estimated number of events during the term of the IHA. The potential total number of individual animals that may be taken is likely an overestimate because the same seal would likely be taken multiple times throughout the season (Table 6).

(7) The anticipated impact of the activity upon the species or stock.

The anticipated impacts of the Estuary management activities are temporary disturbances caused by the presence of staff and equipment, and associated noise, on the beach near the Jenner haulout, and operation of boats and deployment of beach seines near river haulouts. The Water Agency counted seals hauled out and monitored disturbances before, during, and after breaching events from 1996 to 2000 (Merritt-Smith Consulting 1997, 1998, 1999, and 2000; Sonoma County Water Agency and Merritt Smith Consulting 2001). Seals at the Jenner haulout responded most negatively to human disturbances on the beach (typically beach

Table 6. Estimated number of pinnipeds that may be affected (Level B harassment) by Russian River estuary management activities.

Species	No. Animals Expected to Occur	No. Take Events ^a	Potential Total Number of Individual Animals that may be Taken ^{b,c}
Lagoon Outlet Channel Management on the Sandbar (May 15 to October 15)			
Pacific harbor seal	Implementation: 103 ^b Maintenance & Monitoring: May-103; June-100; July-75; Aug-17; Sept-5; Oct-22	Implementation (May-Sept): 3 Maintenance May -1; June-Sept-4/month; Oct-1 Monitoring (June-Sept-2/month; Oct-1	Implementation: 309 Maintenance.: 913 Monitoring: 416 TOTAL: 1,638
California sea lion (potential to encounter once per event May-Oct)	1	3	3
Northern elephant seal (potential to encounter once per event May-Oct)	1	3	3
Artificial Breaching on the Sandbar (October 16 to May 14)			
Pacific harbor seal	Oct: 22 Nov: 11 Dec:134 Jan: 118 Feb: 137 Mar: 167 Apr: 173 May: 103	Oct: 2 ^e Nov: 2 Dec: 2 Jan: 1 Feb: 1 Mar: 1 Apr: 1 May: 1 11 events maximum	Oct: 44 Nov: 22 Dec: 268 Jan: 118 Feb: 137 Mar: 167 Apr: 173 May: 103 TOTAL: 1,032
California sea lion (potential to encounter once per event Oct-Apr)	1	7	7
Northern elephant seal (potential to encounter once per event Dec-May)	1	6	6
Biological and Physical Habitat Monitoring in the Estuary			
Pacific harbor seal	1	65	65
California sea lion (potential to encounter once per month May-Dec)	1	4	4
Northern elephant seal (potential to encounter in Dec)	1	1	1

^a For implementation of the lagoon outlet channel, an event is defined as a single, 2-day episode. It is assumed that the same individual seals would be hauled out during a single event. For the remaining activities, an event is defined as a single day on which an activity occurs. Some events may include multiple activities listed in Table 3.

^b The estimated potential total number of individual animals that may be taken equates to the maximum number of seals of each species anticipated to be encountered per event multiplied by the estimated number of events during the term of the IHA.

^c The potential total number of individual animals that may be taken is likely an overestimate because the same seal would likely be taken multiple times throughout the season

^e The number of events is the monthly average number of artificial breaching events from 1996 to 2010 (Table 1). Based on number of events, the maximum number of artificial breaching events would be 15. The average number of breaching events

from 1996 to 2010 is 6 events/year. The number of events was reduced from the data in Table 1 to 1 for the month of April and 1 for the month of May to anticipate the request from NMFS that only a single closure in each of these months be breached if it seemed unlikely that additional closures would occur prior to May 15th. The potential for two closures in the first two weeks of May seems low. The number of artificial breaching events in February was reduced to one because closures in that month occurred only in 2007, which was an unusual circumstance. December was increased from an average of 1 to 2 events because of the recent experience of multiple closures during heavy storm events.

visitors approaching the haulout and the presence of Water Agency crews and equipment near the haulout). The typical pinniped reactions to disturbances observed were alerts (lifting heads towards source of disturbance), moving to a different location on the beach, or flushing into the water. It is not unusual for pinnipeds to remain on or near the haulout during breaching activities, which may indicate that pinnipeds at the Jenner haulout are somewhat tolerant to disturbance (Heckel 1994).

Stampeding or dead pups have not been observed during monitoring of the Water Agency's artificial breaching activities. Implementation of the lagoon outlet channel, as required by NMFS' Russian River Biological Opinion, has occurred on a single event (July 8, 2010), and the potential direct effects on harbor seals and their pups was similar to artificial breaching activities as construction methods are similar.

More specific data on the behavior of harbor seals during artificial breaching activities, specifically their responses to disturbance, are available in Merritt Smith Consulting (1997, 1998, 1999 and 2001) and Sonoma County Water Agency and Merritt Smith Consulting (2001), and the annual data report for NMFA IHA No. 14426. Mortenson (1996) also discusses harbor seal behavior during the time pups are present. However, water level management activities described in 2009 occurred outside the harbor seal pupping season and so the NMFS IHA No. 14426 data reports does not describe pup behavioral responses to Water Agency activities.

The opportunity for mother/pup bonding at the Jenner haulout is not expected to be impacted by implementation of the lagoon outlet channel or artificial breaching activities. The peak of pupping season is likely by mid-May in most years, and implementation of the lagoon outlet channel would begin around May 15th (as required by the Russian River Biological Opinion). By this time, it is expected that "bonding" between mothers and pups would have likely occurred. The number of artificial breaching activities during the months of March, April and May has been relatively low in the past (see Table 1 of the Water Agency's IHA application), and the breaching activities occur in a single day over several hours. Artificial breaching activities are not expected to impact mother/pup bonding. The proposed project would incorporate the mitigation measures included in the NMFS IHA No. 14426, including limiting access to the beach during the pupping season (March 15 to June 30) as follows:

- If a pup less than one week old is on the beach where heavy machinery would be used or on the path used to access the work location, the breaching event will be delayed until the pup has left the site or the latest day possible to prevent flooding while still maintaining suitable fish rearing habitat. Pups less than one week old should be characterized by being up to 15 kg, thin or their body length, or an umbilicus or natal

pelage is present. The Water Agency will coordinate with the locally established seal monitoring program [SealWatch] to determine if pups less than one week old are on the beach prior to a breaching event;

- A water level management event may not occur for more than two consecutive days unless flooding threats cannot be controlled;
- The Water Agency will maintain a one week (7 day) “no work” period between water level management events (unless flooding is a threat to the low-lying residential community) to allow for adequate disturbance recovery period. During the “no-work” period, equipment must be removed from the beach;
- Physical and biological monitoring, as described in Table 2, will not be conducted if a pup less than one week old is present at the monitoring site or on a path to the site.

Construction of the lagoon outlet channel may require the presence of Water Agency crews and equipment on the beach for up to 2 consecutive days. There have been several breaching events that required up to 2 days of work with a bulldozer or excavator without any apparent long-term impacts to the presence of seals at the haulout. Seals at the Jenner haulout experience regular disturbance by beach visitors and continual noise from the adjacent Highway 1 and would likely only be temporarily disturbed by the presence of Water Agency crews over a 2-day period. However, it is difficult to predict the response to the presence of up to 2 pieces of heavy equipment on the beach during the initial construction of the outlet channel. Monitoring of the pinniped response to this disturbance is included under question 13 below.

During both summer lagoon outlet channel management and artificial breaching activities, Water Agency crews would approach the haulout ahead of the heavy equipment to minimize the potential for flushes to result in a stampede, a particular concern during pupping season. Water Agency staff would avoid walking or driving equipment through the haulout. Crews on foot would take caution to approach the haulout slowly and to make an effort to be seen from a distance, if possible, rather than appearing suddenly at the top of the beach. Seals are usually alerted to the presence of the heavy equipment on the barrier beach well before it approaches the haulout due to the equipment’s noise. Equipment would be driven slowly on the beach and care would be taken to minimize the number of shut downs and start ups when the equipment is on the beach. During the Water Agency’s monitoring from 1996 to 2000, pinnipeds typically abandoned the haulout prior to the bulldozer reaching the breaching location due to disturbance from beach visitors prior to crews arriving onsite. Once breaching was completed, equipment and crews left the beach and pinnipeds returned to the haulout soon after.

(8) The anticipated impact of the activity on the availability of the species or stocks of marine mammals for subsistence uses.

Not applicable.

(9) The anticipated impact of the activity upon the habitat of the marine mammal populations, and the likelihood of restoration of the affected habitat.

The purposes of the lagoon outlet channel management and artificial breaching activities are to manage the sandbar at Goat Rock State Beach to improve summer rearing habitat for juvenile salmonids in the Russian River estuary and to minimize potential flood risk to low-lying properties on the Estuary, respectively. These activities would result in physical alterations of the Jenner haulout. When the barrier beach closes, water surface elevations in the Estuary rise, resulting in the haulout increasing in elevation on the beach, as well as flooding of haulouts in the Russian River. For the summer lagoon outlet channel, elevations would be targeted between 4 and 9 ft NGVD. For artificial breaching activities, the sandbar would be breached when water surface elevations ranged from 4.5 and 7 ft NGVD.

The lagoon outlet channel would alter the beach by creating a shallow outlet channel that would convey river flow to pass over the sandbar and minimize or eliminate tidal exchange from May 15th to October 15th.⁶ The gentle slope of the outlet channel would allow seals to travel through the channel, although the shallow depths (0.5 to 2 ft.) would likely not allow for swimming through the channel. Depending on the barrier beach height and the location of the river's thalweg when the beach closes, part of the outlet channel may be constructed in areas where seals typically haul out on the Estuary side. The outlet channel would be maintained from May 15th to October 15th. After October 15th, the closed barrier beach would be artificially breached when water surface elevations in the Estuary approach 7.0 feet NGVD as read at the Jenner visitor's center. Artificial breaching activities alter the habitat by creating a pilot channel through the closed sandbar. The location of the pilot channel is dependent on the height and width of the sandbar and the location of the river's thalweg. The pilot channel could be constructed in areas where seals typically haul out.

Construction of the lagoon outlet channel and artificial breaching pilot channels requires excavated sand to be side cast on the beach. Any sand excavated would be side cast on the adjacent beach in such a way as to minimize changes to beach topography. During implementation and maintenance of the lagoon outlet channel, side cast sand would be graded on the adjacent beach and/or placed within the wave runup zone on the beach so it is redistributed on the beach. During artificial breaching, the excavated sand is side cast adjacent to the pilot channel because it will be scoured as the mouth opens with flows leaving the estuary.

During the Water Agency's pinniped monitoring associated with artificial breaching activities from 1996 to 2000, the number of harbor seals hauled out at Goat Rock State Beach declined when the barrier beach closed (although the initial decline was often observed to be followed by a short increase in the number of seals at the haulout) and then increased the day following an artificial breaching event (Merritt-Smith Consulting 1997, 1998, 1999, and 2000; Sonoma

⁶ The lagoon management period is May 15th to October 15th, as described in the Russian River Biological Opinion (NMFS 2008).

County Water Agency and Merritt Smith Consulting 2001). This response to barrier beach closure followed by artificial breaching is anticipated to continue. However, less information is available regarding the number of pinnipeds that use the haulout during extended sandbar closure in the lagoon management period (May 15th to October 15th) or when a perched lagoon and outlet channel are established. Collection of baseline information during the lagoon management period would be included in the monitoring described under question 13 below. The Water Agency's previous monitoring, as well as Twohy's daily counts of seals at the sandbar (Table 4) indicate that the number of seals at the haulout declines from August to October, so management of the lagoon outlet channel would have little effect on haulout use. This was the general observation in the fall of 2009 and 2010. The late spring and early summer (May, June, and July), which coincides with pupping season and the likely initiation of lagoon management (following a closure in this time period), may be the most sensitive time period. Pinniped monitoring in 1997 represented some of the longest beach closures in the late spring and early summer months (7 to 11 days). The number of pinnipeds at the haulout varied during the course of the barrier beach closure. Numbers of pinnipeds declined, but then increased, and declined again during the closure and then increased following artificial breaching during each of the spring and early summer sandbar closures in 1997 (Merritt-Smith Consulting 1998). This may indicate that seals present at the haulout during the pupping season are unlikely to completely abandon the haulout from May to July. The outlet channel is anticipated to provide access into the lagoon and to the ocean for pinnipeds at the Jenner haulout. Based on these monitoring results, the numbers of seals hauled out from May through July would be expected to fluctuate, but it is likely that the haulout would continue to be used by harbor seals.

Biological and water quality monitoring would not physically alter pinniped habitat.

(10) The anticipated impact of the loss or modification of the habitat on the marine mammal populations involved.

The modifications of habitat described previously under question 9 would be temporary. The Russian River estuary management activities are anticipated to have minimal effects on the overall habitat of California stocks of Pacific harbor seal and northern elephant seal, or the U.S. stock of California sea lion. Habitat modification effects would be limited to the Jenner haulout at the mouth of the Russian River.

Changes in haulout elevation regularly occur with the tides at this site and any habitat that would be impacted by side cast sand would be temporary. Seals would still have access to the estuary/lagoon waters and could still flush into the water during high water surface elevation periods. Modification of habitat resulting from construction of the lagoon outlet channel or artificial breaching pilot channel would also be temporary in nature. Harbor seals are regularly observed crossing overland from the Pacific Ocean to haul out on the Estuary side of the beach, even in bar-open conditions, so it is anticipated that seals would continue to use the haulout in bar-closed, lagoon conditions.

(11) The availability and feasibility (economic and technological) of equipment, methods, and manner of conducting such activity or other means of effecting the least practicable adverse impact upon the affected species or stocks, their habitat, and on their availability for subsistence uses, paying particular attention to rookeries, mating grounds, and areas of similar significance.

During both summer lagoon outlet channel management and artificial breaching activities, Water Agency crews would approach the haulout ahead of the heavy equipment to minimize the potential for flushes to result in a stampede. Water Agency staff would avoid walking or driving equipment through the haulout. Crews on foot would take caution to approach the haulout slowly and to make an effort to be seen from a distance, if possible, rather than appearing suddenly at the top of the sandbar. Seals are usually alerted to the presence of the heavy equipment on the sandbar well before it approaches the haulout due to the equipment's noise. Equipment would be driven slowly on the beach and care would be taken to minimize the number of shut downs and start ups when the equipment is on the beach to reduce disturbance of seals from loud noises following a relatively quiet period. All work, including monitoring, would be completed as efficiently as possible, with the fewest number of heavy equipment possible, to minimize disturbance of seals at the haulout. Boats operating near river haulouts would be kept within posted speed limits and driven as far from the haulouts as safely possible to minimize flushing seals.

(12) Where the proposed activity would take place in or near a traditional Arctic subsistence hunting area and/or may affect the availability of a species or stock of marine mammal for Arctic subsistence uses, the applicant must submit either a plan of cooperation or information that identifies what measures have been taken and/or would be taken to minimize any adverse effects on the availability of marine mammals for subsistence uses.

Not applicable

(13) The suggested means of accomplishing the necessary monitoring and reporting that would result in increased knowledge of the species, the level of taking or impacts on populations of marine mammals that are expected to be present while conducting activities and suggested means of minimizing burdens by coordinating such reporting requirements with other schemes already applicable to persons conducting such activity. Monitoring plans should include a description of the survey techniques that would be used to determine the movement and activity of marine mammals near the activity site(s) including migration and other habitat uses, such as feeding.

Please see the attached "Russian River Estuary Management Activities Pinniped Monitoring Plan."

(14) Suggested means of learning of, encouraging, and coordinating research opportunities, plans, and activities relating to reducing such incidental taking and evaluating its effects.

All pinniped data collected during the Russian River Estuary management activities at the Russian River would be made available to NMFS, California Department of Parks and Recreation, the Stewards of the Coasts and Redwoods, and the general public.

REFERENCES

Behrens, D. 2008. Inlet Closure and Morphological Behavior in a Northern California Estuary: The Case of the Russian River: University of California, Davis.

California Department of Fish and Game. 1990. California Sea Lion (*Zalophus californianus*). California Habitat Relationship System Online. Available at <http://www.dfg.ca.gov/biogeodata/cwhr/cawildlife.aspx>.

California Department of Fish and Game. 2001. California's Living Marine Resources: A Status Report December 2001

California Department of Fish and Game. 2005. Harbor Seal (*Phoca vitulina*). California Habitat Relationship System Online. Available at <http://www.dfg.ca.gov/biogeodata/cwhr/cawildlife.aspx>.

California Department of Fish and Game. 2009. Northern Elephant Seal. California Wildlife Habitat Relationships System – online. California Department of Fish and Game California Interagency Wildlife Task Group.

Carretta, J. V., K. A. Forney, M. S. Lowry, J. Barlow, J. Baker, B. Hanson, and M. M. Muto. 2007. U.S. Pacific Marine Mammal Stock Assessment Report: 2007. U.S. Department of Commerce, NOAA Technical Memorandum NMFS-SWFSC-414. 320 pp.

Carretta, J., K.A. Forney, M.S. Lowry, J. Barlow, J. Baker, D. Johnston, B. Hanson, M.M. Muto, D. Lynch, and L. Carswell. 2009. NOAA Technical Memorandum NMFS, U.S. Pacific Marine Mammal Stock Assessments: 2008. NOAA-TM-NMFS-SWFSC-434. U.S. Department of Commerce, National Oceanic and Atmospheric Administration, National Marine Fisheries Service, Southwest Fisheries Science Center.

Hanan, D. and M. J. Beeson. 1994. Harbor Seal, *Phoca vitulina richardsi*, Census in California, May-June, 1993. Final Report submitted to NOAA Fisheries/National Marine Fisheries Service, pursuant to Award NA27FX0273-01.

Hanan, D. A. 1996. Dynamics of Abundance and Distribution for Pacific Harbor Seals, *Phoca vitulina richardsi*, on the Coast of California. Ph.D. Dissertation, University of California, Los Angeles:158 pp.

- Hanson, L.C. 1993. The foraging ecology of harbor seals, *Phoca vitulina*, and California sea lions, *Zalophus californianus*, at the mouth of the Russian River, California. Masters thesis, Sonoma State University. May 9, 1993.
- Heckel, M. 1994. Russian River Estuary Study 1992-1993. Prepared for Sonoma County Department of Planning and California State Coastal Conservancy. .186 pp.
- Herder, M. J. 1986. Seasonal movements and hauling site fidelity of harbor seals, *Phoca vitulina richardsi*, tagged at the Russian River, California. MS. Thesis. Humboldt State University:52 pp.
- Huber, H.R., A.C. Rovetta, L.A. Fry, and S. Johnston. 1991. Age specific natality of northern elephant seals at the South Farallon Islands, California. J. Mammalogy. 72: 525-534.
- Le Boeuf, B. J., D. Crocker, S. Blackwell, and P. Morris. 1993. Sex differences in diving and foraging behaviour of northern elephant seals. In: I. Boyd (ed.). Marine Mammal: Advances in Behavioural and Population Biology. Oxford Univ. Press.
- Lowry, M. S., J. V. Carretta, and K. A. Forney. 2005. Pacific Harbor Seal, *Phoca vitulina richardsi*, Census in California During May-July 2004. NOAA National Marine Fisheries Service Southwest Fisheries Science Center Technical Memorandum SWFSC-Administrative Report LJ-05-06:38 pp.
- Merritt Smith Consulting. 1997. Biological and Water Quality Monitoring in the Russian River Estuary, 1996. Prepared for Sonoma County Water Agency. February 21, 1997.
- Merritt Smith Consulting. 1998. Biological and Water Quality Monitoring in the Russian River Estuary, 1997. Second Annual Report. Prepared for the Sonoma County Water Agency. February 5, 1998.
- Merritt Smith Consulting. 1999. Biological and Water Quality Monitoring in the Russian River Estuary, 1998. Third Annual Report. Prepared for the Sonoma County Water Agency. March 15, 1999.
- Merritt Smith Consulting. 2000. Biological and Water Quality Monitoring in the Russian River Estuary, 1999. Fourth Annual Report. Prepared for the Sonoma County Water Agency. March 24, 2000.
- Mortenson, J. 1996. Human interference with harbor seals at Jenner, CA, 1994-1995. Stewards of Slavianka, Sonoma Coast State Beaches, Russian River/Mendocino Park District. July 11, 1996.
- Mortenson, J. 2009. Location of pinniped haulouts at the Russian River Estuary. April 20, 2009.
- Mortenson, J. and M. Follis. 1997. Northern elephant seal (*Mirounga angustirostris*) aggression on harbor seal (*Phoca vitulina*) pups. Marine Mammal Science 13(3): 526-530.

Mortenson, J. and E. Twohy. 1994. Harbor seals at Jenner, CA, 1974-1993. Stewards of Slavianka, California Department of Parks and Recreation. Duncans Mills, CA.:35 pp.

NMFS (National Marine Fisheries Service). 2008. Russian River Biological Opinion. September 24, 2008.

PWA (Philip Williams and Associates, Ltd.). 2009. Russian River Estuary Outlet Channel Adaptive Management Plan Year 1, Draft. Prepared for Sonoma County Water Agency Philip Williams & Associates, Ltd. with Bodega Marine Laboratory, University Of California at Davis. PWA Ref. # 1958.01. May 1, 2009.

Sonoma County Water Agency and Merritt Smith Consulting. 2001. Biological and Water Quality Monitoring in the Russian River Estuary, 2000. Fifth Annual Report. June 12, 2001.

Stewart, B., B. Le Boeuf, P. Yochem, H. Huber, R. DeLong, R. Jameson, W. Sydeman, and S. Allen. 1994. History and present status of the northern elephant seal population. In: B.J. Le Boeuf and R.W. Laws (eds.) Elephant Seals. Univ. Calif. Press, Berkeley. 414 pp.

Stewart, B.S. and H.R. Huber. 1993. Mammalian species: *Mirounga angustirostris*. *American Journal of Mammalogists* 449:1-10.