

## SPERM WHALE (*Physeter macrocephalus*): Northern Gulf of Mexico Stock

### STOCK DEFINITION AND GEOGRAPHIC RANGE

Sperm whales are found throughout the world's oceans in deep waters to the edge of the ice at both poles (Leatherwood and Reeves 1983; Rice 1989; Whitehead 2002). Sperm whales were commercially hunted in the Gulf of Mexico by American whalers from sailing vessels until the early 1900s (Townsend 1935). In the northern Gulf of Mexico (i.e., U.S. Gulf of Mexico) systematic aerial and ship surveys indicate that sperm whales inhabit continental slope and oceanic waters where they are widely distributed (Figure 1; Fulling *et al.* 2003; Mullin and Fulling 2004; Mullin *et al.* 2004; Maze-Foley and Mullin 2006; Mullin 2007). Seasonal aerial surveys confirm that sperm whales are present in the northern Gulf of Mexico in all seasons (Mullin *et al.* 1994; Hansen *et al.* 1996; Mullin and Hoggard 2000). The information for southern Gulf of Mexico waters is more limited, but there are sighting and stranding records from each season with sightings widely distributed in continental slope waters of the western Bay of Campeche (Ortega-Ortiz 2002).

Sperm whales throughout the world exhibit a geographic social structure where females and juveniles of both sexes occur in mixed groups and inhabit tropical and subtropical waters. Males, as they mature, initially form bachelor groups but eventually become more socially isolated and more wide-ranging, inhabiting temperate and polar waters as well (Whitehead 2003). While this pattern also applies to the Gulf of Mexico, results of multi-disciplinary research conducted in the Gulf since 2000 confirms speculation by Schmidly

(1981) and indicates clearly that Gulf of Mexico sperm whales constitute a stock that is distinct from other Atlantic Ocean stocks(s) (Mullin *et al.* 2003; Jaquet 2006; Jochens *et al.* 2008). The following summarizes the most significant stock structure-related findings from Jochens *et al.* (2008). Measurements of the total length of Gulf of Mexico sperm whales indicate that they are 1.5-2.0m smaller on average compared to whales measured in other areas. Female/immature group size in the Gulf (9-11 whales) is about one-third that found elsewhere. Tracks from 39 whales satellite tagged in the northern Gulf were monitored for up to 607 days. No discernable seasonal migrations were made, but Gulf-wide movements primarily along the northern Gulf slope did occur. The tracks showed that whales exhibit a range of movement patterns within the Gulf, including movement into the southern Gulf in a few cases, but that only 1 whale (a male) left the Gulf of Mexico. This animal moved into the North Atlantic and then back into the Gulf after about 2 months. Additionally, no matches were found when 285 individual whales photo-identified from the Gulf and about 2500 from the North Atlantic and Mediterranean were compared. An analysis of matrilineally inherited mtDNA revealed a significant genetic differentiation between animals from the northern Gulf of Mexico compared to those from the North Atlantic Ocean, North Sea and Mediterranean Sea. Analysis of biparentally inherited nuclear DNA showed no significant difference between whales sampled in the Gulf and those from the other areas of the North Atlantic, indicating that mature males move in and out of the Gulf. Sperm whales make vocalizations used in a social context called "codas" that have distinct patterns that are apparently culturally transmitted, and based on degree of social affiliation, mixed groups of sperm whales




Figure 1. Distribution of sperm whale sightings from SEFSC spring vessel surveys during 1996-2001 and from summer 2003 and spring 2004 surveys. All the on-effort sightings are shown, though not all were used to estimate abundance. Solid lines indicate the 100m and 1,000m isobaths and the offshore extent of the U.S. EEZ.

worldwide can be placed in recognizable acoustic clans. Recordings from mixed groups in the Gulf of Mexico compared to those from other areas of the Atlantic indicated that Gulf sperm whales constitute a distinct acoustic clan that is rarely encountered outside of the Gulf. It is assumed from this that groups from other clans enter the northern Gulf only infrequently.

## POPULATION SIZE

The best abundance estimate available for northern Gulf of Mexico sperm whales is 1,665 (CV=0.20) (Mullin 2007; Table 1). This estimate is pooled from summer 2003 and spring 2004 oceanic surveys covering waters from the 200-m isobath to the seaward extent of the U.S. Exclusive Economic Zone (EEZ).

### Earlier abundance estimates

Estimates of abundance were derived through the application of distance sampling analysis (Buckland *et al.* 2001) and the computer program DISTANCE (Thomas *et al.* 1998) to sighting data. From 1991 through 1994, line-transect vessel surveys were conducted in conjunction with bluefin tuna ichthyoplankton surveys during spring in the northern Gulf of Mexico from the 200-m isobath to the seaward extent of the U.S. EEZ (Hansen *et al.* 1995). Annual cetacean surveys were conducted along a fixed plankton sampling trackline. Survey effort-weighted estimated average abundance of sperm whales for all surveys combined was 530 (CV=0.31) (Hansen *et al.* 1995; Table 1). Similar surveys were conducted during spring from 1996 to 2001 (excluding 1998) in oceanic waters of the northern Gulf of Mexico. Due to limited survey effort in any given year, survey effort was pooled across all years to develop an average abundance estimate. The estimate of abundance for sperm whales in oceanic waters, pooled from 1996 to 2001, is 1,349 (CV=0.23) (Mullin and Fulling 2004; Table 1).

### Recent surveys and abundance estimates

During summer 2003 and spring 2004, line-transect surveys dedicated to estimating the abundance of oceanic cetaceans were conducted in the northern Gulf of Mexico. During each year, a grid of uniformly-spaced transect lines from a random start were surveyed from the 200-m isobath to the seaward extent of the U.S. EEZ using NOAA Ship *Gordon Gunter* (Mullin 2007).

As recommended in the GAMMS Workshop Report (Wade and Angliss 1997), estimates older than 8 years are deemed unreliable, and therefore should not be used for PBR determinations. Because most of the data for estimates prior to 2003 were older than this 8-year limit and due to the different sampling strategies, estimates from the 2003 and 2004 surveys were considered most reliable. The estimate of abundance for sperm whales in oceanic waters, pooled from 2003 to 2004, was 1,665 (CV=0.20) (Mullin 2007; Table 1), which is the best available abundance estimate for this species in the northern Gulf of Mexico.

Month/Year	Area	$N_{best}$	CV
Apr-Jun 1991-1994	Oceanic waters	530	0.31
Apr-Jun 1996-2001 (excluding 1998)	Oceanic waters	1,349	0.23
Jun-Aug 2003, Apr-Jun 2004	Oceanic waters	1,665	0.20

### Minimum Population Estimate

The minimum population estimate is the lower limit of the two-tailed 60% confidence interval of the log-normal distributed abundance estimate. This is equivalent to the 20th percentile of the log-normal distributed abundance estimate as specified by Wade and Angliss (1997). The best estimate of abundance for sperm whales is 1,665 (CV=0.20). The minimum population estimate for the northern Gulf of Mexico is 1,409 sperm whales.

### Current Population Trend

There are insufficient data to determine the population trends for this species. The pooled abundance estimate for 2003-2004 of 1,665 (CV=0.20) and that for 1996-2001 of 1,349 (CV=0.29) are not significantly different ( $P>0.05$ ), but due to the precision of the estimates, the power to detect a difference is relatively low. These estimates are 2-3 times larger than that for 1991-1994 of 530 (CV=0.31). The 2003-2004 estimates were based on less negatively biased estimates of sperm whale group size and may account for part of the difference. Nevertheless, these temporal abundance estimates are difficult to interpret without a Gulf of Mexico-wide understanding of sperm

whale abundance. The Gulf of Mexico is composed of waters belonging to the U.S., Mexico and Cuba. U.S. waters only comprise about 40% of the entire Gulf of Mexico, and 65% of oceanic waters are south of the U.S. EEZ. The oceanography of the Gulf of Mexico is quite dynamic, and the spatial scale of the Gulf is small relative to the ability of most cetacean species to travel. Studies based on abundance and distribution surveys restricted to U.S. waters are unable to detect temporal shifts in distribution beyond U.S. waters that might account for any changes in abundance.

### **CURRENT AND MAXIMUM NET PRODUCTIVITY RATES**

Current and maximum net productivity rates are unknown for this stock. For purposes of this assessment, the maximum net productivity rate was assumed to be 0.04. This value is based on theoretical modeling showing that cetacean populations may not grow at rates much greater than 4% given the constraints of their reproductive history (Barlow *et al.* 1995).

### **POTENTIAL BIOLOGICAL REMOVAL**

Potential biological removal level (PBR) is the product of the minimum population size, one half the maximum net productivity rate and a recovery factor (MMPA Sec. 3.16 U.S.C. 1362; Wade and Angliss 1997). The minimum population size is 1,409. The maximum productivity rate is 0.04, the default value for cetaceans. The “recovery” factor, which accounts for endangered, depleted, threatened stocks, or stocks of unknown status relative to optimum sustainable population (OSP), is assumed to be 0.1 because the sperm whale is an endangered species. PBR for the northern Gulf of Mexico sperm whale is 2.8.

### **ANNUAL HUMAN-CAUSED MORTALITY AND SERIOUS INJURY**

There has been no reported fishing-related mortality of a sperm whale during 1998-2007 (Yeung 1999; 2001; Garrison 2003; Garrison and Richards 2004; Garrison 2005; Fairfield Walsh and Garrison 2006; Fairfield Walsh and Garrison 2007; Fairfield and Garrison 2008).

### **Fisheries Information**

The level of past or current, direct, human-caused mortality of sperm whales in the northern Gulf of Mexico is unknown. Pelagic swordfish, tunas and billfish are the targets of the longline fishery operating in the northern Gulf of Mexico. There were no reports of mortality or serious injury to sperm whales by this fishery.

A commercial fishery for sperm whales operated in the Gulf of Mexico in deep waters between the Mississippi River delta and DeSoto Canyon during the late 1700s to the early 1900s (Mullin *et al.* 1991), but the exact number of whales taken is not known (Townsend 1935; Lowery 1974). Townsend (1935) reported many records of sperm whales from April through July in the north-central Gulf (Petersen and Hoggard 1996).

### **Other Mortality**

Two sperm whale strandings were documented during 2007 (1 in Florida, 1 in Texas). No sperm whale strandings were documented during 2004-2006. A total of 9 sperm whale strandings were documented in the northern Gulf of Mexico during 1999-2003 (Table 2; NOAA National Marine Mammal Health and Stranding Response Database unpublished data, accessed 16 September 2008). No evidence of human interactions was detected for these stranded animals. Stranding data probably underestimate the extent of fishery-related mortality and serious injury because not all of the marine mammals which die or are seriously injured in fishery interactions wash ashore, not all that wash ashore are discovered, reported or investigated, nor will all of those that do wash ashore necessarily show signs of entanglement or other fishery interaction. Finally, the level of technical expertise among stranding network personnel varies widely as does the ability to recognize signs of fishery interactions.

Seismic vessel operations in the Gulf of Mexico (commercial and academic) now operate with marine mammal observers as part of required mitigation measures. There have been no reported seismic-related or industry ship-related mortalities or injuries to sperm whales. However, disturbance by anthropogenic noise may prove to be an important habitat issue in some areas of this population’s range, notably in areas of oil and gas activities and/or where shipping activity is high. Results from very limited studies of northern Gulf of Mexico sperm whale responses to seismic exploration indicate that sperm whales do not appear to exhibit horizontal avoidance of seismic survey activities. Data did suggest that there may be some decrease in foraging effort during exposure to full-array airgun firing, at least for some individuals. Further study is needed as samples sizes are insufficient at this time (Jochens *et al.* 2008).

The potential impact, if any, of coastal pollution may be an issue for this species in portions of its habitat, though little is known on this to date.

Table 2. Sperm whale (*Physeter macrocephalus*) strandings along the northern Gulf of Mexico coast, 1999-2003. No sperm whale strandings were documented during 2004-2006.

STATE	1999	2000	2001	2002	2003	TOTAL
Alabama	0	0	0	0	0	0
Florida	1	2	1	1	1	6
Louisiana	1	0	0	0	1	2
Mississippi	0	0	0	0	0	0
Texas	0	1	0	0	0	1
<b>TOTAL</b>	2	3	1	1	2	9

### STATUS OF STOCK

The status of sperm whales in the northern Gulf of Mexico, relative to OSP, is unknown. This species is listed as endangered under the Endangered Species Act (ESA). There are insufficient data to determine the population trends for this species. Total human-caused mortality and serious injury for this stock is not known. There is insufficient information available to determine whether the total fishery-related mortality and serious injury for this stock is insignificant and approaching zero mortality and serious injury rate. This is a strategic stock because the sperm whale is listed as an endangered species under the ESA.

### REFERENCES CITED

- Barlow, J., S.L. Swartz, T.C. Eagle and P.R. Wade 1995. U.S. marine mammal stock assessments: Guidelines for preparation, background, and a summary of the 1995 assessments. NOAA Tech. Memo. NMFS-OPR-6. 73 pp.
- Buckland, S.T., D.R. Anderson, K.P. Burnham, J.L. Laake, D.L. Borchers and L. Thomas 2001. Introduction to distance sampling: estimating abundance of biological populations. Oxford University Press. 432 pp.
- Fairfield, C.P. and L.P. Garrison 2008. Estimated bycatch of marine mammals and sea turtles in the US Atlantic pelagic longline fleet during 2007. NOAA Tech. Memo. NOAA NMFS-SEFSC-572. 62 pp.
- Fairfield Walsh, C. and L.P. Garrison 2006. Estimated bycatch of marine mammals and turtles in the U.S. Atlantic pelagic longline fleet during 2005. NOAA Tech. Memo. NOAA NMFS-SEFSC-539. 52 pp.
- Fairfield Walsh, C. and L.P. Garrison 2007. Estimated bycatch of marine mammals and turtles in the U.S. Atlantic pelagic longline fleet during 2006. NOAA Tech. Memo. NOAA NMFS-SEFSC-560. 54 pp.
- Fulling, G.L., K.D. Mullin and C.W. Hubard 2003. Abundance and distribution of cetaceans in outer continental shelf waters of the U.S. Gulf of Mexico. Fish. Bull. 101: 923-932.
- Garrison, L.P. 2003. Estimated bycatch of marine mammals and turtles in the U.S. Atlantic pelagic longline fleet during 2001-2002. NOAA Tech. Memo. NMFS-SEFSC-515. 52 pp.
- Garrison, L.P. 2005. Estimated bycatch of marine mammals and turtles in the U.S. Atlantic pelagic longline fleet during 2004. NOAA Tech. Memo. NMFS-SEFSC-531. 57 pp.
- Garrison, L.P. and P.M. Richards 2004. Estimated bycatch of marine mammals and turtles in the U.S. Atlantic pelagic longline fleet during 2003. NOAA Tech. Memo. NMFS-SEFSC-527. 57 pp.
- Hansen, L.J., K.D. Mullin, T.A. Jefferson and G.P. Scott 1996. Visual surveys aboard ships and aircraft. Pages 55-132 in: R.W. Davis and G.S. Fargion, (eds.) Distribution and abundance of marine mammals in the north-central and western Gulf of Mexico: Final report. Volume II: Technical report. OCS Study MMS 96-0027. Minerals Management Service, Gulf of Mexico OCS Region, New Orleans, LA. .
- Hansen, L.J., K.D. Mullin and C.L. Roden 1995. Estimates of cetacean abundance in the northern Gulf of Mexico from vessel surveys. Southeast Fisheries Science Center, Miami, FL. Contribution No. MIA-94/95-25. 9 pp.
- Jaquet, N. 2006. A simple photogrammetric technique to measure sperm whales at sea. Mar. Mamm. Sci. 22(4): 862-879.
- Jochens, A., D. Biggs, K. Benoit-Bird, D. Engelhaupt, J. Gordon, C. Hu, N. Jaquet, M. Johnson, R. Leben, B. Mate, P. Miller, J. Ortega-Ortiz, A. Thode, P. Tyack and B. Würsig 2008. Sperm whale seismic study in the Gulf

- of Mexico: Synthesis report. U.S. Dept. of the Interior, Minerals Management Service, Gulf of Mexico OCS Region, New Orleans, LA. OCS Study MMS 2008-006. 341 pp.
- Leatherwood, S. and R.R. Reeves 1983. The Sierra Club handbook of whales and dolphins. Sierra Club Books, San Francisco. 302 pp.
- Lowery, G.H., Jr. 1974. The mammals of Louisiana and its adjacent waters. Louisiana State University Press, Baton Rouge. 565 pp.
- Maze-Foley, K. and K.D. Mullin 2006. Cetaceans of the oceanic northern Gulf of Mexico: Distributions, group sizes and interspecific associations. *J. Cetacean Res. Manage.* 8(2): 203-213.
- Mullin, K., W. Hoggard, C. Roden, R. Lohofener, C. Rogers and B. Taggart 1991. Cetaceans on the upper continental slope in the north-central Gulf of Mexico. OCS Study/MMS 91-0027. U.S. Dep. Interior, Minerals Management Service, Gulf of Mexico OCS Regional Office, New Orleans, LA 108 pp.
- Mullin, K., W. Hoggard, C. Roden, R. Lohofener, C. Rogers and B. Taggart 1994. Cetaceans on the upper continental slope in the north-central Gulf of Mexico. *Fish. Bull.* 92: 773-786.
- Mullin, K.D. 2007. Abundance of cetaceans in the oceanic Gulf of Mexico based on 2003-2004 ship surveys. Available from: NMFS, Southeast Fisheries Science Center, P.O. Drawer 1207, Pascagoula, MS 39568, 26 pp.
- Mullin, K.D., D. Engelhaupt, C.E. Cates and N.B. Barros 2003. Sperm whale research in the Gulf of Mexico. International Whaling Commission Working Paper SC/55/O15. Available from: NMFS, Southeast Fisheries Science Center, P.O. Drawer 1207, Pascagoula, MS 39568.
- Mullin, K.D. and G.L. Fulling 2004. Abundance of cetaceans in the oceanic northern Gulf of Mexico. *Mar. Mamm. Sci.* 20(4): 787-807.
- Mullin, K.D. and W. Hoggard 2000. Visual surveys of cetaceans and sea turtles from aircraft and ships. Pages 111-172 in: R.W. Davis, W.E. Evans and B. Würsig, (eds.) *Cetaceans, sea turtles and seabirds in the northern Gulf of Mexico: Distribution, abundance and habitat associations. Volume II: Technical report.* Minerals Management Service, Gulf of Mexico OCS Region, New Orleans. OCS Study MMS 96-0027.
- Mullin, K.D., W. Hoggard and L.J. Hansen 2004. Abundance and seasonal occurrence of cetaceans in outer continental shelf and slope waters of the north-central and northwestern Gulf of Mexico. *Gulf of Mexico Science* 2004(1): 62-73.
- Ortega-Ortiz, J.G. 2002. Multiscale analysis of cetacean distribution in the Gulf of Mexico. Ph.D. thesis. Texas A&M University. 170 pp.
- Petersen, J.C. and W. Hoggard 1996. First sperm whale (*Physeter macrocephalus*) record in Mississippi. *Gulf Research Reports* 9(3): 215-217.
- Rice, D.W. 1989. Sperm whale, *Physeter macrocephalus* Linnaeus, 1758. Pages 177-233 in: S.H. Ridgway and R. Harrison, (eds.) *Handbook of marine mammals, Vol. 4: River dolphins and the larger toothed whales.* Academic Press, London.
- Schmidly, D.J. 1981. Marine mammals of the southeastern United States and the Gulf of Mexico. U.S. Fish and Wildlife Service, Office of Biological Services, Washington, DC. FWS/OBS-80/41. 165 pp.
- Thomas, L., J.L. Laake, J.F. Derry, S.T. Buckland, D.L. Borchers, D.R. Anderson, K.P. Burnham, S. Strindberg, S.L. Hedley, F.F.C. Marques, J.H. Pollard and R.M. Fewster 1998. Distance 3.5. Research Unit for Wildlife Population Assessment, University of St. Andrews, St. Andrews, UK.
- Townsend, C.H. 1935. The distribution of certain whales as shown by logbook records of American whale ships. *Zoologica* 19: 1-50.
- Wade, P.R. and R.P. Angliss 1997. Guidelines for assessing marine mammal stocks: Report of the GAMMS Workshop April 3-5, 1996, Seattle, Washington. NOAA Tech. Memo. NMFS-OPR-12. 93 pp.
- Whitehead, H. 2002. Estimates of the current global population size and historical trajectory for sperm whales. *Mar. Ecol. Prog. Ser.* 242: 295-304.
- Whitehead, H. 2003. Sperm whales: Social evolution in the ocean. The University of Chicago Press, Chicago, IL. 431 pp.
- Yeung, C. 1999. Estimates of marine mammal and marine turtle bycatch by the U.S. Atlantic pelagic longline fleet in 1998. NOAA Tech. Memo. NMFS-SEFSC-430. 26 pp.
- Yeung, C. 2001. Estimates of marine mammal and marine turtle bycatch by the U.S. Atlantic pelagic longline fleet in 1999-2000. NOAA Tech. Memo., NMFS, Southeast Fisheries Science Center, Miami, FL. NMFS-SEFSC-467. 43 pp.

