

COMMON BOTTLENOSE DOLPHIN (*Tursiops truncatus truncatus*): Western North Atlantic Central Florida Coastal Stock

STOCK DEFINITION AND GEOGRAPHIC RANGE

Geographic Range and Coastal Morphotype Habitat

In the western North Atlantic the coastal morphotype of common bottlenose dolphins is continuously distributed in nearshore coastal and estuarine waters along the U.S. Atlantic coast south of Long Island, New York, around the Florida peninsula and into the Gulf of Mexico. Based on differences in mitochondrial DNA haplotype frequencies, coastal animals in the northern Gulf of Mexico and the western North Atlantic represent separate stocks (Duffield and Wells 2002; Rosel *et al.* 2009).

The coastal morphotype is morphologically and genetically distinct from the larger, more robust morphotype that occupies habitats further offshore in the western North Atlantic and Gulf of Mexico (Mead and Potter 1995; Hoelzel *et al.* 1998; Rosel *et al.* 2009; Vollmer 2011). Aerial surveys conducted between 1978 and 1982 (Kenney 1990) north of Cape Hatteras, North Carolina identified two concentrations of bottlenose dolphins, one near the coast within the 25-m isobath and the other further offshore concentrated at the continental shelf edge. The lowest density of bottlenose dolphins was observed over the continental shelf. It was suggested, therefore, that north of Cape Hatteras, North Carolina, the coastal morphotype is restricted to waters <25 m deep (Kenney 1990). Similar patterns were observed during summer months in more recent aerial surveys (Garrison and Yeung 2001; Garrison *et al.* 2003). However, south of Cape Hatteras during both winter and summer months, there was no clear longitudinal discontinuity in bottlenose dolphin sightings (Garrison and Yeung 2001; Garrison *et al.* 2003).

To address the question of the distribution of coastal and offshore morphotypes in waters south of Cape Hatteras, tissue samples were collected in coastal, shelf and slope waters from New England to Florida between 1997 and 2006. Genetic analyses using mitochondrial DNA sequences of these biopsies identified individual animals to the coastal or offshore morphotype. Using the genetic results from all surveys combined, a logistic regression was used to model the probability that a particular bottlenose dolphin group was of the coastal morphotype as a function of environmental variables including depth, sea surface temperature and distance from shore. These models were used to partition the bottlenose dolphin groups observed during aerial surveys between the two morphotypes (Garrison *et al.* 2003).

The genetic results and spatial patterns observed in aerial surveys indicate both regional and seasonal differences in the longitudinal distribution of the two morphotypes in coastal Atlantic waters. During summer months, all biopsy samples collected from coastal waters north of Cape Lookout, North Carolina (<20 m deep) were of the coastal morphotype, and all samples collected in deeper waters (>40 m deep) were of the offshore morphotype. South of Cape Lookout, the probability of an observed bottlenose dolphin group being of the coastal morphotype declined with increasing depth. In intermediate depth waters, there was spatial overlap between the two morphotypes. Offshore morphotype bottlenose dolphins were observed at depths as shallow as 13 m, and coastal morphotype dolphins were observed at depths of 31 m and 75 km from shore (Garrison *et al.* 2003).

Winter samples were collected primarily from coastal waters in North Carolina and Georgia and the vast majority of them were of the coastal morphotype; however, one offshore morphotype group was sampled during November just south of Cape Lookout only 7.3 km from shore. Coastal morphotype samples were also collected farther away from shore at 33 m depth and 39 km from shore. The logistic regression model for this region indicated a decline in the probability of a coastal morphotype group with increasing distance from shore; however, the model predictions were highly uncertain due to limited sample sizes and spatial overlap between the two morphotypes. Samples collected in Georgia waters also indicated significant overlap between the two morphotypes with a declining probability of the coastal morphotype with increasing depth. A coastal morphotype sample was collected 112 km from shore at a depth of 38 m. An offshore sample was collected in 22 m depth at 40 km from shore. As with the North Carolina model, the Georgia logistic regression predictions are uncertain due to limited sample size and high overlap between the two morphotypes (Garrison *et al.* 2003).

In summary, the primary habitat of the coastal morphotype of bottlenose dolphin in the western North Atlantic extends from Florida to Long Island, New York, during summer months and in waters less than 20 m deep, including estuarine and inshore waters. South of Cape Lookout, the coastal morphotype also occurs in lower densities over the continental shelf and overlaps spatially with the offshore morphotype.

Distinction between Coastal and Estuarine Bottlenose Dolphins

In addition to inhabiting coastal nearshore waters, the coastal morphotype of common bottlenose dolphin also inhabits inshore estuarine waters along the U.S. east coast and Gulf of Mexico (Wells *et al.* 1987; Scott *et al.* 1990; Wells *et al.* 1996; Weller 1998; Zolman 2002; Speakman *et al.* 2006; Stolen *et al.* 2007; Balmer *et al.* 2008; Mazzoil *et al.* 2008). There are multiple lines of evidence supporting demographic separation between bottlenose dolphins residing within different estuaries along the Atlantic coast. For example, long-term photo-identification (photo-ID) studies in waters around Charleston, South Carolina, have identified communities of resident dolphins that are seen within relatively restricted home ranges year-round (Zolman 2002; Speakman *et al.* 2006). In Biscayne Bay, Florida, there is a similar community of bottlenose dolphins with evidence of year-round residents that are genetically distinct from animals residing in a nearby estuary in Florida Bay (Litz *et al.* 2012). A long-term photo-ID study in the Indian River Lagoon system in central Florida has also identified year-round resident dolphins repeatedly observed across multiple years (Stolen *et al.* 2007; Mazzoil *et al.* 2008). A few published studies demonstrate that these resident animals are genetically distinct from animals in nearby coastal waters. A study conducted near Jacksonville, Florida demonstrated significant genetic differences between animals in coastal and estuarine waters (Caldwell 2001; Rosel *et al.* 2009) and animals resident in the Charleston estuarine system show significant genetic differentiation from animals biopsied in coastal waters of southern Georgia (Rosel *et al.* 2009).

Despite evidence for genetic differentiation between estuarine and coastal populations, the degree of spatial overlap between these populations remains unclear. Photo-ID studies within estuaries demonstrate seasonal immigration and emigration and the presence of transient animals (e.g., Speakman *et al.* 2006). In addition, the degree of movement of resident estuarine animals into coastal waters on seasonal or shorter time scales is poorly understood. Bottlenose dolphins inhabiting primarily estuarine habitats are considered distinct stocks from those inhabiting coastal habitats. Bottlenose dolphin stocks inhabiting coastal waters are the focus of this report.

Definition of the Central Florida Coastal Stock

Initially, a single stock of common bottlenose dolphins of the coastal morphotype was thought to migrate seasonally between New Jersey (summer months) and central Florida based on seasonal patterns in strandings during a large scale mortality event occurring during 1987-1988 (Scott *et al.* 1988). However, re-analysis of stranding data (McLellan *et al.* 2003) and extensive analysis of genetic (Rosel *et al.* 2009), photo-ID (Zolman 2002) and satellite telemetry (Hohn and Hansen, NMFS unpublished data) data demonstrate a complex mosaic of coastal bottlenose dolphin stocks. Integrated analysis of these multiple lines of evidence suggests that there are five coastal stocks of bottlenose dolphins: the Northern Migratory and Southern Migratory Stocks, a South Carolina/Georgia Coastal Stock, a Northern Florida Coastal Stock and a Central Florida Coastal Stock.

The spatial extent of these stocks, their potential seasonal movements, and their relationships with estuarine stocks are poorly understood. Migratory movement and spatial distribution of the Northern Migratory Stock is best understood based on tag-telemetry, photo-ID and aerial survey data and migrates seasonally between coastal waters of central North Carolina and New Jersey. It is not thought to overlap with the Central Florida Coastal Stock in any season. The Southern Migratory Coastal Stock is defined primarily on satellite tag telemetry studies and is thought to migrate south from waters of southern Virginia and north central North Carolina in the summer to waters south of Cape Fear and as far south as coastal Florida during winter months. It is unclear whether this stock overlaps with the Central Florida Coastal Stock in any season.

During summer months when the Southern Migratory Coastal Stock is found in waters north of Cape Fear, North Carolina, bottlenose dolphins are still seen in coastal waters of South Carolina, Georgia and Florida, indicating the presence of additional stocks of coastal animals. Speakman *et al.* (2006) using photo-ID studies documented dolphins in coastal waters off Charleston, South Carolina, that are not known resident members of the estuarine stock. Genetic analyses of samples from northern Florida, Georgia and central South Carolina (primarily the estuaries around Charleston), using both mitochondrial DNA and nuclear microsatellite markers indicate significant genetic differences between these areas (NMFS 2001; Rosel *et al.* 2009). This stock assessment report addresses the Central Florida Coastal stock, which is present in coastal Atlantic waters from 29.4°N south to the western end of Vaca Key (~24.69°N -81.11°W) where the stock boundary for the Florida Keys Stock begins (Figure 1). There has been little study of bottlenose dolphin stock structure in coastal waters of southern Florida, therefore the southern boundary of the Central Florida Stock is uncertain. There is no obvious boundary defining the offshore extent of this stock. The combined genetic and logistic regression analysis (Garrison *et al.* 2003) indicated that in waters less than 10 m depth, 70% of the bottlenose dolphins were of the coastal morphotype. Between 10 and 20 m depth, the percentage of animals of the coastal morphotype dropped precipitously, and at depths >40 m nearly all (>90%) animals were of the offshore morphotype. These spatial patterns may not apply in the Central Florida Coastal Stock, as there is a significant change in the bathymetric slope and a close approach of the Gulf Stream to the shoreline south of Cape Canaveral.

POPULATION SIZE

The best available estimate for the Central Florida Coastal Stock of common bottlenose dolphins in the western North Atlantic is 4,895 (CV=0.71; Table 1). This estimate is from aerial surveys conducted during the summers of 2010 and 2011 covering waters from Florida to New Jersey.

Earlier abundance estimates

Earlier abundance estimates for the Central Florida Coastal Stock were derived from aerial surveys conducted during the summer of 2002 and 2004. Survey tracklines were set perpendicular to the shoreline and included coastal waters to depths of 40 m. These surveys employed two observer teams operating independently on the same aircraft to estimate visibility bias. In summer 2004, an additional aerial survey between central Florida and New Jersey was conducted. As with the 2002 surveys, effort was stratified into 0-20 m and 20-40 m strata with the majority of effort in the shallow depth stratum. Observed common bottlenose dolphin groups from these were partitioned between the coastal and offshore morphotypes based upon analysis of available biopsy samples (Garrison *et al.* 2003). The previous best abundance estimate was based upon an average of the estimates from the 2002 and 2004 aerial surveys. This estimate was 6,318 (CV=0.26).




Figure 1. The Central Florida Coastal stock of common bottlenose dolphins (29.4°N to Vaca Key). Symbols represent all sightings of bottlenose dolphin groups from NMFS 2010 & 2011 aerial surveys; dark symbols- groups within the boundaries of this stock. In waters >20 m, sightings may include the offshore morphotype of bottlenose dolphins.

Recent surveys and abundance estimates

The Southeast Fisheries Science Center conducted aerial surveys of continental shelf waters along the U.S. East Coast from southeastern Florida to Cape May, New Jersey, during the summers of 2010 and 2011. The surveys were conducted along tracklines oriented perpendicular to the shoreline that were latitudinally spaced 20 km apart and covered waters from the shoreline to the continental shelf break. The summer 2010 survey was conducted during 24 July–14 August 2010, and 7,944 km of on-effort tracklines completed. A total of 127 common bottlenose dolphin groups were observed including 1,541 animals. During the 2011 summer survey, 8,665 km of trackline were completed between Cape May, New Jersey and Ft. Pierce, Florida. The survey was conducted during 6 July - 29 July 2011. The 2011 survey also included more closely spaced “fine-scale” tracklines in waters offshore of New Jersey and Virginia within areas being evaluated for the placement of offshore energy installations. A total of 112 bottlenose dolphin groups were sighted including 1,339 animals.

Both the summer 2010 and 2011 surveys were conducted using a two-team approach to develop estimates of visibility bias using the independent observer approach with Distance analysis (Laake and Borchers 2004). However, the detection functions from both surveys indicated a decreased probability of detection near the trackline, which limited the effectiveness of the method for correcting for visibility bias due to a relatively small number of sightings made by both teams near the trackline. Abundance estimates were therefore derived by combining the sightings from both teams during a survey and “left-truncating” the data by analyzing only sightings occurring greater than 100 m from the trackline during the 2010 survey and 50 m during the 2011 survey (Buckland *et al.* 2001). Detection functions were fit to these left-truncated data accounting for the effects of survey conditions (e.g., sea state, glare, water color) on the detection probabilities. A logistic regression model was used to estimate the probability that a given group of dolphins observed during the aerial survey was of the coastal vs. offshore morphotype as a function of water depth (Garrison *et al.* 2003). This probability was incorporated into the abundance estimation to derive an estimate of coastal morphotype dolphins observed during the 2010 and 2011 aerial surveys. A bootstrap resampling approach was used to estimate the variance of the estimates. The resulting abundance estimates assume that detection probability at the truncation distance is equal to 1. While the estimates could not be explicitly corrected for this assumption, analyses of the summer 2010 data suggest that this bias is likely small.

The abundance estimates for the Central Florida Coastal Stock were based upon tracklines and sightings occurring between 29.4°N latitude and Ft. Pierce, Florida and in waters from the shoreline to the 40-m isobath. The abundance estimate derived from the summer 2010 survey was 9,842 (CV=0.84), and the estimate from the summer 2011 survey was 1,338 (CV=0.65). The best estimate is a weighted average of these two with higher weighting given to the more precise estimate from 2011. The resulting best estimate is 4,895 (CV=0.71).

| Table 1. Summary of abundance estimates for the western North Atlantic Central Florida Coastal Stock of common bottlenose dolphins. Month, year, and area covered during each abundance survey, and resulting abundance estimate (N_{best}) and coefficient of variation (CV). | | | |
|--|-----------------------------------|------------|------|
| Month/Year | Area | N_{best} | CV |
| Summer 2002 and 2004 | 29.4°N Latitude to Ft. Pierce, FL | 6,318 | 0.26 |
| Summer 2010 and 2011 | 29.4°N Latitude to Ft. Pierce, FL | 4,895 | 0.71 |

Minimum Population Estimate

The minimum population size (N_{min}) for each stock was calculated as the lower bound of the 60% confidence interval for a log-normally distributed mean (Wade and Angliss 1997). The best estimate for the Central Florida Coastal Stock is 4,895 (CV=0.71). The resulting minimum population estimate is 2,851.

Current Population Trend

A trend analysis has not been conducted for this stock. There are 2 estimates from the 2002/2004 (6,318; CV=0.26) and 2010/2011 (4,895; CV=0.71) surveys. Methodological differences between the estimates need to be evaluated to quantify trends.

CURRENT AND MAXIMUM NET PRODUCTIVITY RATES

Current and maximum net productivity rates are not known for the western North Atlantic coastal morphotype. The maximum net productivity rate was assumed to be 0.04. This value is based on theoretical modeling showing that cetacean populations may not grow at rates much greater than 4% given the constraints of their reproductive life history (Barlow *et al.* 1995).

POTENTIAL BIOLOGICAL REMOVAL

Potential Biological Removal (PBR) is the product of the minimum population size, one-half the maximum productivity rate, and a “recovery” factor (MMPA Sec. 3. 16 U.S.C. 1362; Wade and Angliss 1997). The minimum population size of the Central Florida Coastal Stock of common bottlenose dolphins is 2,851. The maximum productivity rate is 0.04, the default value for cetaceans. The recovery factor is 0.5 because this stock is depleted. PBR for this stock of common bottlenose dolphins is 29.

ANNUAL HUMAN-CAUSED MORTALITY AND SERIOUS INJURY

The total annual human-caused mortality and serious injury for the Central Florida Coastal Stock during 2009–2013 is unknown because this stock is known to interact with unobserved fisheries (see below). The mean annual fishery-related mortality and serious injury for strandings identified as fishery-caused was 0.2. No additional mortality or serious injury was documented from other human-caused actions. The minimum total mean annual human-caused mortality and serious injury for this stock during 2009–2013 was 0.2.

Fishery Information

The commercial fisheries that interact, or that potentially could interact, with this stock are the Category II Southeastern U.S. Atlantic shark gillnet; Southeast Atlantic gillnet; Atlantic blue crab trap/pot; and Southeastern U.S. Atlantic, Gulf of Mexico stone crab trap/pot fisheries; and the Category III Florida spiny lobster trap/pot; and Atlantic Ocean, Gulf of Mexico, Caribbean commercial passenger fishing vessel (hook and line) fisheries. Only limited observer data are available for these and other fisheries that may interact with this stock. Detailed fishery information is presented in Appendix III.

Southeastern U.S. Atlantic Shark Gillnet Fishery and Southeast Atlantic Gillnet

Gillnet fisheries targeting finfish and sharks operate in southeast waters between North Carolina and southern Florida. There have been no observed bottlenose dolphin takes within the stock boundaries since 2003. Historically, a drift net fishery targeting coastal sharks operated in waters including within the Central Florida Coastal Stock boundaries during winter months. Common bottlenose dolphin takes ($n=2$) were observed in the drift net fisheries targeting sharks in 2002 and 2003 (Garrison 2007). Currently, gillnet fisheries include a number of different fishing methods and gear types including drift nets, “strike” fishing, and anchored (“sink”) gillnets. The majority of this fishing is reported from waters of North Carolina and central Florida. However, there has been a significant reduction in the amount of drift gillnet fishing targeting sharks during the last several years.

Trap/Pot

During 2009–2013, 1 stranded animal assigned to the Central Florida Coastal Stock was reported entangled in trap/pot gear. The animal was disentangled and released alive seriously injured (Maze-Foley and Garrison in prep). It was not possible to determine which specific trap/pot fishery (blue crab, stone crab or spiny lobster) interacted with this stranding. Because there is no systematic observer program, it is not possible to estimate the total number of interactions or mortalities associated with trap/pot gear.

Hook and Line

During 2009–2013, no interactions were documented involving hook and line gear entanglement or ingestion within the Central Florida Coastal Stock area. Earlier interactions have been documented. It is not possible to estimate the total number of interactions with hook and line gear because there is no systematic observer program.

Other Mortality

During 2009–2013, 137 stranded common bottlenose dolphins were recovered in the waters of the Central Florida Coastal Stock (NOAA National Marine Mammal Health and Stranding Response Database unpublished data, accessed 11 June 2014). It could not be determined if there was evidence of human interaction for 90 of these strandings, and for 45 it was determined there was no evidence of human interaction. The remaining 2 showed evidence of human interactions, both of which were fisheries interactions. As mentioned above, 1 animal was

reported entangled in trap/pot fishery gear and was released alive seriously injured (Maze-Foley and Garrison in prep). Stranding data probably underestimate the extent of human and fishery-related mortality and serious injury because not all of the dolphins that die or are seriously injured in human interactions wash ashore, or, if they do, they are not all recovered (Peltier *et al.* 2012; Wells *et al.* 2015). Additionally, not all carcasses will show evidence of human interaction, entanglement or other fishery-related interaction due to decomposition, scavenger damage, etc. (Byrd *et al.* 2014). Finally, the level of technical expertise among stranding network personnel varies widely as does the ability to recognize signs of human interaction.

The Indian River Lagoon Estuarine System (IRLES) Stock experienced an Unusual Mortality Event (UME) in 2008. From May to August a total of 47 bottlenose dolphins from the IRLES Stock and 1 dolphin from the Central Florida Coastal Stock were considered to be part of this UME (NOAA National Marine Mammal Health and Stranding Response Database unpublished data, accessed 13 September 2012). Infectious disease is suspected as a possible cause of this event. A UME was declared in the summer of 2013 for the mid-Atlantic coast from New York to Brevard County, Florida. Beginning in July 2013, bottlenose dolphins have been stranding at elevated rates. The total number of stranded bottlenose dolphins from New York through North Florida (Brevard County) as of mid-October 2014 (1 July 2013 - 19 October 2014) was ~1546. Morbillivirus has been determined to be the cause of the event. Most strandings and morbillivirus positive animals have been recovered from the ocean side beaches rather than from within the estuaries, suggesting that at least so far coastal stocks have been more impacted by this UME than estuarine stocks. However, the UME is still ongoing as of December 2014 when this report was drafted, and work continues to determine the effect of this event on all bottlenose dolphin stocks in the Atlantic.

HABITAT ISSUES

The nearshore and estuarine habitats occupied by the coastal morphotype are adjacent to areas of high human population and some are highly industrialized. The blubber of stranded dolphins examined during the 1987-1988 mortality event contained very high concentrations of organic pollutants (Kuehl *et al.* 1991). More recent studies have examined persistent organic pollutant concentrations in bottlenose dolphin inhabiting estuaries along the Atlantic coast and have likewise found evidence of high blubber concentrations particularly near Charleston, South Carolina, and Beaufort, North Carolina (Hansen *et al.* 2004). The concentrations found in male dolphins from both of these sites exceeded toxic threshold values that may result in adverse effects on health or reproductive rates (Schwacke *et al.* 2002; Hansen *et al.* 2004). Studies of contaminant concentrations relative to life history parameters showed higher levels of mortality in first-born offspring and higher contaminant concentrations in these calves and in primiparous females (Wells *et al.* 2005). The exposure to environmental pollutants and subsequent effects on population health is an area of concern and active research.

STATUS OF STOCK

Common bottlenose dolphins are not listed as threatened or endangered under the Endangered Species Act, but the Central Florida Coastal Stock is a strategic stock due to the depleted listing under the Marine Mammal Protection Act. From 1995 to 2001, NMFS recognized only a single migratory stock of coastal bottlenose dolphins in the western North Atlantic, and the entire stock was listed as depleted. This stock structure was revised in 2002 to recognize both multiple stocks and seasonal management units and again in 2008 and 2010 to recognize resident estuarine stocks and migratory and resident coastal stocks. This stock retains the depleted designation as a result of its origins from the originally delineated depleted coastal migratory stock. PBR for the Central Florida Coastal Stock is 29 and so the zero mortality rate goal, 10% of PBR, is 2.9. The documented mean annual human-caused mortality for this stock for 2009 – 2013 was 0.2. However, there are several commercial fisheries operating within this stock's boundaries and these fisheries have little to no observer coverage. Therefore, the documented mortality must be considered a minimum estimate of total fishery-related mortality. Insufficient information is available to determine whether the total fishery-related mortality and serious injury for this stock is insignificant and approaching a zero mortality and serious injury rate. The status of this stock relative to OSP is unknown. There are insufficient data to determine population trends for this stock.

REFERENCES CITED

Balmer, B.C., R.S. Wells, S.M. Nowacek, D.P. Nowacek, L.H. Schwacke, W.A. McLellan, F.S. Scharf, T.K. Rowles, L.J. Hansen, T.R. Spradlin and D.A. Pabst. 2008. Seasonal abundance and distribution patterns of common bottlenose dolphins (*Tursiops truncatus*) near St. Joseph Bay, Florida, USA. *J. Cetacean Res. Manage.* 10: 157-167.

- Barlow, J., S.L. Swartz, T.C. Eagle and P.R. Wade. 1995. U.S. marine mammal stock assessments: Guidelines for preparation, background and a summary of the 1995 assessments. NOAA Tech. Memo. NMFS-OPR-6, 73 pp.
- Buckland, S. T., D. R. Andersen, K. P. Burnham, J. L. Laake, D. L. Borchers and L. Thomas. 2001. Introduction to distance sampling: Estimating abundance of biological populations. Oxford University Press, New York, 432 pp.
- Byrd, B.L., A.A. Hohn, G.N. Lovewell, K.M. Altman, S.G. Barco, A. Friedlaender, C.A. Harms, W.A. McLellan, K.T. Moore, P.E. Rosel and V.G. Thayer. 2014. Strandings illustrate marine mammal biodiversity and human impacts off the coast of North Carolina, USA. Fish. Bull. 112: 1-23.
- Caldwell, M. 2001. Social and genetic structure of bottlenose dolphin (*Tursiops truncatus*) in Jacksonville, Florida. Ph.D. dissertation from University of Miami. 143 pp.
- Duffield, D.A. and R.S. Wells. 2002. The molecular profile of a resident community of bottlenose dolphins, *Tursiops truncatus*. pp. 3-11. In: C. J. Pfeiffer, (ed.) Cell and Molecular Biology of Marine Mammals. Krieger Publishing, Melbourne, FL. 464 pp.
- Garrison, L.P. 2007. Estimated marine mammal and turtle bycatch in the shark gillnet fisheries along the southeast U.S. Atlantic coast: 2000-2006. Southeast Fisheries Science Center, 75 Virginia Beach Dr., Miami, FL 33149. PRD Document #07/08-10, 22 pp.
- Garrison, L.P., P.E. Rosel, A.A. Hohn, R. Baird and W. Hoggard. 2003. Abundance of the coastal morphotype of bottlenose dolphin *Tursiops truncatus*, in U.S. continental shelf waters between New Jersey and Florida during winter and summer 2002. NMFS/SEFSC report prepared and reviewed for the Bottlenose Dolphin Take Reduction Team. Available from: NMFS, Southeast Fisheries Science Center, 75 Virginia Beach Dr., Miami, FL 33149.
- Garrison, L.P. and C. Yeung. 2001. Abundance estimates for Atlantic bottlenose dolphin stocks during summer and winter, 1995. NMFS/SEFSC report prepared and reviewed for the Bottlenose Dolphin Take Reduction Team. Available from: NMFS, Southeast Fisheries Science Center, 75 Virginia Beach Dr., Miami, FL 33149.
- Hansen, L.J., L.H. Schwacke, G.B. Mitchum, A.A. Hohn, R.S. Wells, E.S. Zolman and P.A. Fair. 2004. Geographic variation in polychlorinated biphenyl and organohaline pesticide concentrations in the blubber of bottlenose dolphins from the US Atlantic coast. Sci. Total Environ. 319: 147-172.
- Hoelzel, A.R., C.W. Potter, and P.B. Best. 1998. Genetic differentiation between parapatric nearshore and offshore populations of the bottlenose dolphin. Proc. Royal Soc. London 265: 1177-1183.
- Kenney, R.D. 1990. Bottlenose dolphins off the northeastern United States. pp. 369-386. In: S. Leatherwood and R. Reeves (eds.) The bottlenose dolphin. Academic Press, San Diego, CA. 653 pp.
- Kuehl, D.W., R. Haebler, C. Potter. 1991. Chemical residues in dolphins from the US Atlantic coast including Atlantic bottlenose obtained during the 1987/1988 mass mortality. Chemosphere 22: 1071-1084.
- Laake, J.L. and D.L. Borchers. 2004. Methods for incomplete detection at distance zero. pp. 108-189. In: S.T. Buckland, D.R. Andersen, K.P. Burnham, J.L. Laake and L. Thomas (eds.) Advanced distance sampling. Oxford University Press, New York.
- Litz, J.A., C.R. Hughes, L.P. Garrison, L.A. Fieber and P.E. Rosel. 2012. Genetic structure of common bottlenose dolphins (*Tursiops truncatus*) inhabiting adjacent South Florida estuaries - Biscayne Bay and Florida Bay. J. Cetacean Res. Manage. 12(1): 107-117.
- Maze-Foley, K. and L.P. Garrison. in prep. Preliminary serious injury determinations for small cetaceans off the southeast U.S. coast, 2007-2011.
- Mazzoil, M., J.S. Reif, M. Youngbluth, M.E. Murdoch, S.E. Bechdel, E. Howells, S.D. McCulloch, L.J. Hansen and G.D. Bossart. 2008. Home ranges of bottlenose dolphins (*Tursiops truncatus*) in the Indian River Lagoon, Florida: Environmental correlates and implications for management strategies. EcoHealth 5(3): 278-288.
- Mead, J.G. and C.W. Potter. 1995. Recognizing two populations of the bottlenose dolphin (*Tursiops truncatus*) off the Atlantic coast of North America: Morphological and ecological considerations. IBI Reports 5: 31-44.
- McLellan, W.M., A.S. Friedlaender, J.G. Mead, C.W. Potter and D.A. Pabst. 2003. Analysing 25 years of bottlenose dolphin (*Tursiops truncatus*) strandings along the Atlantic coast of the USA: Do historic records support the coastal migratory stock hypothesis? J. Cetacean Res. Manage. 4: 297-304.
- NMFS. 2001. Stock structure of coastal bottlenose dolphins along the Atlantic coast of the US. NMFS/SEFSC Report prepared for the Bottlenose Dolphin Take Reduction Team. Available from: NMFS, Southeast Fisheries Science Center, 75 Virginia Beach Dr., Miami, FL 33149.
- Peltier, H., W. Dabin, P. Daniel, O. Van Canneyt, G. Dorémus, M. Huon and V. Ridoux. 2012. The significance of stranding data as indicators of cetacean populations at sea: modelling the drift of cetacean carcasses. Ecol.

Indicators 18: 278–290.

- Rosel, P.E., L. Hansen and A.A. Hohn. 2009. Restricted dispersal in a continuously distributed marine species: Common bottlenose dolphins *Tursiops truncatus* in coastal waters of the western North Atlantic. *Mol. Ecol.* 18: 5030–5045.
- Schwacke, L.H., E.O. Voit, L.J. Hansen, R.S. Wells, G.B. Mitchum, A.A. Hohn and P.A. Fair. 2002. Probabilistic risk assessment of reproductive effects of polychlorinated biphenyls on bottlenose dolphins (*Tursiops truncatus*) from the southeast United States coast. *Env. Toxic. Chem.* 21: 2752-2764.
- Scott, G.P., D.M. Burn and L.J. Hansen. 1988. The dolphin die off: Long term effects and recovery of the population. *Proceedings: Oceans '88, IEEE Cat. No. 88-CH2585-8, Vol. 3: 819-823.*
- Scott, M.D., R.S. Wells and A.B. Irvine. 1990. A long-term study of bottlenose dolphins on the west coast of Florida. pp. 235-244. *In: S. Leatherwood and R. R. Reeves (eds.) The bottlenose dolphin. Academic Press, San Diego, CA. 653 pp.*
- Stolen, M.K., W.N. Durden and D.K. Odell. 2007. Historical synthesis of bottlenose dolphin (*Tursiops truncatus*) stranding data in the Indian River Lagoon system, Florida, from 1977-2005. *Fla. Sci.* 70: 45-54.
- Speakman, T., E.S. Zolman, J. Adams, R.H. Defran, D. Laska, L. Schwacke, J. Craigie and P. Fair. 2006. Temporal and spatial aspects of bottlenose dolphin occurrence in coastal and estuarine waters near Charleston, South Carolina. NOAA Tech. Memo. NOS-NCCOS-37, 243 pp.
- Vollmer, N.L. 2011. Population structure of common bottlenose dolphins in coastal and offshore waters of the Gulf of Mexico revealed by genetic and environmental analyses. Ph.D. Dissertation from University of Louisiana at Lafayette. 420 pp.
- Wade, P.R. and R.P. Angliss. 1997. Guidelines for assessing marine mammal stocks: Report of the GAMMS Workshop April 3-5, Seattle, Washington. NOAA Tech. Memo. NMFS-OPR-12, 93 pp.
- Weller, D.W. 1998. Global and regional variation in the biology and behavior of bottlenose dolphins. Ph.D. dissertation from Texas A&M University, College Station. 142 pp.
- Wells, R.S., J.B. Allen, G. Lovewell, J. Gorzelany, R.E. Delynn, D.A. Fauquier and N.B. Barros. 2015. Carcass-recovery rates for resident bottlenose dolphins in Sarasota Bay, Florida. *Mar. Mamm. Sci.* 31(1): 355-368.
- Wells, R.S., M.D. Scott and A.B. Irvine. 1987. The social structure of free ranging bottlenose dolphins. pp. 247-305. *In: H. Genoways (ed.) Current Mammalogy, Vol. 1. Plenum Press, New York. 519 pp.*
- Wells, R.S., V. Tornero, A. Borrell, A. Aguilar, T.K. Rowles, H.L. Rhinehart, S. Hofmann, W.M. Jarman, A.A. Hohn and J.C. Sweeney. 2005. Integrating life history and reproductive success data to examine potential relationships with organochlorine compounds for bottlenose dolphins (*Tursiops truncatus*) in Sarasota Bay, Florida. *Sci. Total Environ.* 349: 106-119.
- Wells, R.S., K.W. Urian, A.J. Read, M.K. Bassos, W.J. Carr and M.D. Scott. 1996. Low-level monitoring of bottlenose dolphins, *Tursiops truncatus*, in Tampa Bay, Florida: 1988-1993. NOAA Tech. Memo. NMFS-SEFSC- 385, 25 pp. + 6 Tables, 8 Figures, and 4 Appendices.
- Zolman, E. S. 2002. Residence patterns of bottlenose dolphins (*Tursiops truncatus*) in the Stono River Estuary, Charleston County, South Carolina. *Mar. Mamm. Sci.* 18: 879-892.

