

## **BRYDE'S WHALE (*Balaenoptera edeni*): Eastern Tropical Pacific Stock**

### **STOCK DEFINITION AND GEOGRAPHIC RANGE**

The International Whaling Commission (IWC) recognizes 3 stocks of Bryde's whales in the North Pacific (eastern, western, and East China Sea), 3 stocks in the South Pacific (eastern, western and Solomon Islands), and one cross-equatorial stock (Peruvian) (Donovan 1991). Bryde's whales are distributed widely across the tropical and warm-temperate Pacific (Leatherwood et al. 1982), and there is no justification for splitting stocks between the northern and southern hemispheres (Donovan 1991). Past surveys have shown them to be common and distributed throughout the eastern tropical Pacific with a concentration around the equator east of 110°W (corresponding approximately to the IWC's "Peruvian stock") and a lower densities west of 140°W (Lee 1993; Wade and Gerrodette 1993). They are also the most common baleen whale in the central Gulf of California (Tershy et al. 1990). Sightings and acoustic recordings of Bryde's whales in southern California waters have increased in the past decade (Kerosky et al. 2012, Smultea et al. 2012), possibly signaling a northward range expansion (Kerosky et al. 2012). Acoustic recordings indicate Bryde's whales are present in southern California waters from summer through early winter (Kerosky et al. 2012). At least seven sightings have been documented in southern / central California waters between 1991 and 2014 (Barlow and Forney 2007, Smultea et al. 2012, Barlow 2016). Bryde's whales in California waters likely belong to a larger population inhabiting at least the eastern part of the tropical Pacific. Acoustic call types of Bryde's whales in southern California waters match a type found along the west coast of Baja California (Kerosky et al. 2012). For the Marine Mammal Protection Act (MMPA) stock assessment reports, Bryde's whales within the Pacific U.S. Exclusive Economic Zone are divided into two areas: 1) the eastern tropical Pacific (east of 150°W and including the Gulf of California and waters off California; this report), and 2) Hawaiian waters.

### **POPULATION SIZE**

In the western North Pacific, Bryde's whale abundance in the early 1980s was estimated independently by tag mark-recapture and ship survey methods to be 22,000 to 24,000 (Tillman and Mizroch 1982; Miyashita 1986). Bryde's whale abundance has never been estimated for the entire eastern Pacific; however, a portion of that stock in the eastern tropical Pacific was estimated as 13,000 (CV=0.20; 95% CI = 8,900-19,900) (Wade and Gerrodette 1993), and the minimum number in the Gulf of California was estimated at 160 based on individually-identified whales (Tershy et al. 1990). The most recent verified sighting in California waters occurred in 2014 during a systematic line-transect survey designed to estimate cetacean abundance (Barlow 2016). That sighting did not occur during standard search effort and thus, no estimate of abundance is available from the 2014 survey.

### **Minimum Population Estimate**

The only minimum estimate of Bryde's whale abundance for the eastern tropical Pacific (11,163; Wade and Gerrodette 1993) is over 8 years old and thus, no current estimate of minimum abundance is available.

### **Current Population Trend**

There are no data on trends in Bryde's whale abundance in the eastern tropical Pacific.

### **CURRENT AND MAXIMUM NET PRODUCTIVITY RATES**

There are no estimates of the growth rate of Bryde's whale populations in the Pacific (Best 1993).

### **POTENTIAL BIOLOGICAL REMOVAL**

The potential biological removal (PBR) level for this stock cannot be calculated because a current abundance estimate is unavailable.

### **HUMAN CAUSED MORTALITY**

#### **Historic Whaling**

The reported take of North Pacific Bryde's whales by commercial whalers totaled 15,076 in the western Pacific from 1946-1983 (Holt 1986) and 2,873 in the eastern Pacific from 1973-81 (Cooke 1983). In addition, 2,304 sei-or-Bryde's whales were taken in the eastern Pacific from 1968-72 (Cooke 1983) (based on subsequent catches, most of these were probably Bryde's whales). None were reported taken by shore-based whaling stations in central or northern California between 1919 and 1926 (Clapham et al. 1997) or 1958 and 1965 (Rice 1974). There has been a prohibition on taking Bryde's whales since 1988.

**Table 1.** Summary of available information on the incidental mortality and injury of Bryde's whales (eastern tropical Pacific stock) for commercial fisheries that might take this species (Carretta *et al.* 2014a, 2012a, 2012b, Carretta and Enriquez 2009, 2010; Carretta *et al.* 2004). n/a indicates that data are not available. Mean annual takes are based on 2001-2013 data unless noted otherwise.

Fishery Name	Year(s)	Data Type	Percent Observer Coverage	Observed mortality (and injury in parentheses)	Estimated mortality (CV in parentheses)	Mean annual takes (CV in parentheses)
CA/OR thresher shark/swordfish drift gillnet fishery	2001-2013	observer	19%	0	0	0
<b>Total annual takes</b>						0

### Fishery Information

The California swordfish drift gillnet fishery is the only fishery that is likely to take Bryde's whales from this stock, but no entanglements have been observed (Table 1). Detailed information on this fishery is provided in Appendix 1. Mean annual takes for this fishery are zero (Table 1) and are based on 2001-2013 data, the period during which a season/area closure has limited most fishing to southern California waters. Although Bryde's whales have not been observed entangled in California gillnets, some gillnet mortality of large whales may go unobserved because whales swim away with a portion of the net.

### Ship Strikes

One Bryde's whale was documented to have been killed by a ship strike in 2010 (Carretta et al. 2014b, Carretta et al. 2015). The whale was initially sighted alive in Washington state waters with propeller marks and stranded dead about a week later. The mean annual serious injury and mortality rate of Bryde's whales over the most recent 5-year period (2009-2013) is 0.2 whales annually.

### STATUS OF STOCK

Commercial whaling of Bryde's whales was largely limited to the western Pacific. Bryde's whales are not listed as "threatened" or "endangered" under the Endangered Species Act (ESA). Bryde's whales in the eastern tropical Pacific would not be considered a strategic stock under the MMPA. The total human-caused mortality rate is 0.2 whales annually. Current abundance of this stock is unknown and therefore PBR cannot be calculated for this stock. Likewise, human-caused mortality cannot be evaluated in the context of PBR. Increasing levels of anthropogenic sound in the world's oceans has been suggested to be a habitat concern for whales, particularly for baleen whales that may communicate using low-frequency sound.

### REFERENCES

- Barlow, J., and Forney, K.A. 2007. Abundance and population density of cetaceans in the California Current ecosystem. *Fishery Bulletin*, 105(4), 509-526.
- Barlow, J. 2016. Cetacean abundance in the California Current estimated from ship-based line-transect surveys in 1991-2014. Draft Document PSRG-2016-06 presented to the Pacific Scientific Review Group, 25-26 February 2016, Seattle, WA.
- Best, P. B. 1993. Increase rates in severely depleted stocks of baleen whales. *ICES J. Mar. Sci.* 50:169-186.
- Carretta, J.V., M.M. Muto, S. Wilkin, J. Greenman, K. Wilkinson, M. DeAngelis, J. Viezbicke, D. Lawson, J. Rusin, and J. Jannot. 2015. Sources of human-related injury and mortality for U.S. Pacific west coast marine mammal stock assessments, 2009-2013. U.S. Department of Commerce, NOAA Technical Memorandum, NOAA-TM-NMFS-SWFSC-548. 108 p.

- Carretta, J.V., L. Enriquez, and Charles Villafana. 2014a. Marine mammal, sea turtle, and seabird bycatch in California gillnet fisheries in 2012. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-SWFSC-526, 16 p.
- Carretta, J.V., S.M. Wilkin, M.M. Muto, K. Wilkinson, and J. Rusin. 2014b. Sources of human-related injury and mortality for U.S. Pacific west coast marine mammal stock assessments, 2008-2012. NOAA Technical Memorandum, NOAA-TM-NMFS-SWFSC-533. 110 p.
- Carretta, J.V. and L. Enriquez. 2012a. Marine mammal and seabird bycatch in California gillnet fisheries in 2011. NOAA Technical Memorandum, NOAA-TM-NMFS-SWFSC-500. 14p.
- Carretta, J.V. and L. Enriquez. 2012b. Marine mammal and seabird bycatch in California gillnet fisheries in 2010. Administrative Report LJ-12-01. NOAA Fisheries, Southwest Fisheries Science Center, National Marine Fisheries Service, La Jolla, CA 92037. 14 p.
- Carretta, J.V. and L. Enriquez. 2010. Marine mammal and sea turtle bycatch in the California/Oregon swordfish and thresher shark drift gillnet fishery in 2009. Administrative Report LJ-10-03, available from Southwest Fisheries Science Center, 3333 North Torrey Pines Ct., La Jolla, California, 92037. 11p.
- Carretta, J.V. and L. Enriquez. 2009. Marine mammal bycatch in the California/Oregon swordfish and thresher shark drift gillnet fishery in 2008. Administrative Report LJ-09-03, available from Southwest Fisheries Science Center, 3333 North Torrey Pines Ct., La Jolla, California, 92037. 10 p.
- Carretta, J.V., T. Price, D. Petersen, and R. Read. 2004. Estimates of marine mammal, sea turtle, and seabird mortality in the California drift gillnet fishery for swordfish and thresher shark, 1996-2002. *Marine Fisheries Review* 66(2):21-30.
- Clapham, P. J., S. Leatherwood, I. Szczepaniak, and R. L. Brownell, Jr. 1997. Catches of humpback and other whales from shore stations at Moss Landing and Trinidad, California, 1919-1926. *Mar. Mamm. Sci.* 13(3):368-394.
- Cooke, J. G. 1983. Estimates of the stock of Bryde's whales fished off the coast of Peru. *Rept. Int. Whal. Commn.* 33:453-456.
- Donovan, G. P. 1991. A review of IWC stock boundaries. *Rept. Int. Whal. Commn., Special Issue* 13:39-68.
- Holt, S. 1986. Aspects of the assessment and regulation of Bryde's whales in the Northwest Pacific. *Rept. Int. Whal. Commn.* 36:257-262.
- Kerosky, S. M., Širović, A., Roche, L. K., Baumann-Pickering, S., Wiggins, S. M., & Hildebrand, J. A. 2012. Bryde's whale seasonal range expansion and increasing presence in the Southern California Bight from 2000 to 2010. *Deep Sea Research Part I: Oceanographic Research Papers*, 65, 125-132.
- Leatherwood, S., R. R. Reeves, W. F. Perrin, and W. E. Evans. 1982. Whales, dolphins, and porpoises of the eastern North Pacific and adjacent Arctic waters: A guide to their identification. NOAA Technical Rept. NMFS Circular 444. 245 pp.
- Lee, T. 1993. Summary of cetacean survey data collected between the years of 1974 and 1985. NOAA Tech. Memo. NOAA-TM-NMFS-SWFSC-181. 184 pp.
- Miyashita, T. 1986. Sighting estimate for the Bryde's whale stock in the western North Pacific. *Rept. Int. Whal. Commn.* 36:249-252.
- Rice, D. W. 1974. Whales and whale research in the eastern North Pacific. pp. 170-195 *In*: W. E. Schevill (ed.). *The Whale Problem: A Status Report*. Harvard Press, Cambridge, MA.
- Smultea, M. A., Douglas, A. B., Bacon, C. E., Jefferson, T. A., & Mazzuca, L. 2012. Bryde's whale (*Balaenoptera brydei/edeni*) sightings in the Southern California Bight. *Aquatic Mammals*, 38(1), 92-97.
- Tershy, B. R., D. Breese, and C. S. Strong. 1990. Abundance, seasonal distribution and population composition of balaenopterid whales in the Canal de Ballenas, Gulf of California, Mexico. *Rept. Int. Whal. Commn., Special Issue* 12:369-375.
- Tillman, M. F. and S. A. Mizroch. 1982. Mark-recapture estimates of abundance for the western Northern Pacific stock of Bryde's whales. *Rept. Int. Whal. Commn.* 32:335-337.
- Wade, P. R. and T. Gerrodette. 1993. Estimates of cetacean abundance and distribution in the eastern tropical Pacific. *Rept. Int. Whal. Commn.* 43:477-493.