

barndoor skate




Figure A1. Northeast Region barndoor skate has a controlled fishing mortality and biomass is rebuilding. B_{msy} proxy is in kg/tow.

*Skate stocks F/F_{msy} are measured by percent change over a 3 year period.

bluefish (except Gulf of Mexico)




Figure A2. Northeast Region bluefish (except Gulf of Mexico) has a controlled fishing mortality and biomass is rebuilding.

bocaccio




Figure A3. Northwest Region bocaccio has a controlled fishing mortality and biomass is rebuilding as expected.

canary rockfish




Figure A4. Northwest Region canary rockfish has a controlled fishing mortality and biomass is rebuilding as expected.




Figure A5. Northwest Region cowcod has a controlled fishing mortality and biomass is rebuilding as expected.

darkblotched rockfish




Figure A6. Northwest Region darkblotched rockfish has a controlled fishing mortality and biomass is rebuilding as expected.

golden tilefish




Figure A7. Northeast Region golden tilefish has a controlled fishing mortality and biomass is rebuilding.

haddock - Georges Bank




Figure A8. Northeast Region haddock – Georges Bank has a controlled fishing mortality and biomass is rebuilding.

haddock - Gulf of Maine




Figure A9. Northeast Region haddock – Gulf of Maine has a controlled fishing mortality and biomass has increased following the overfished declaration. B_{msy} proxy is in kg/tow.

king mackerel - GoM




Figure A10. South Atlantic Region king mackerel – Gulf of Mexico has a controlled fishing mortality and biomass is rebuilding as expected. B_{msy} proxy is in trillions of eggs.

Pacific ocean perch




Figure A11. Northwest Region Pacific ocean perch has a controlled fishing mortality and biomass is rebuilding as expected. *Declared overfished in 1999 B/B_{msy} was assumed < 0.5 .




Figure A12. Northeast Region pollock has a controlled fishing mortality and biomass is rebuilding. B_{msy} proxy is in kg/tow.




Figure A13. South Atlantic Region red porgy has a controlled fishing mortality and biomass is rebuilding as expected.

redfish




Figure A14. Northeast Region redfish has a controlled fishing mortality and biomass is rebuilding.




Figure A15. Northeast Region spiny dogfish has a controlled fishing mortality and biomass is rebuilding. B_{msy} proxy is in female biomass.




Figure A16. Highly Migratory Species swordfish - North Atlantic has a controlled fishing mortality and biomass is rebuilding. FMP not internationally implemented

$F/F_{msy}^* = 2.19$ for initial
overfished declaration

widow rockfish




Figure A17. Northwest Region widow rockfish has a controlled fishing mortality and biomass is rebuilding as expected.




Figure A18. Northwest Region yelloweye rockfish has a controlled fishing mortality and biomass is rebuilding as expected.